

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/53, 15/55, 15/60, C12P 7/20, 7/18, C12N 1/15, 1/19, 1/21, 9/04, 9/16, 9/88</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/28480</b> <b>(43) International Publication Date:</b> 10 June 1999 (10.06.99)
<b>(21) International Application Number:</b> PCT/US98/25551 <b>(22) International Filing Date:</b> 2 December 1998 (02.12.98)  <b>(30) Priority Data:</b> 08/982,783                      2 December 1997 (02.12.97)                      US  <b>(71) Applicants (for all designated States except US):</b> E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). GENECOR INTERNATIONAL, INC. [US/US]; 4 Cambridge Place, 1870 Winton Road, Rochester, NY 14618 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> NAIR, Ramesh, V. [IN/US]; 905-D Cloister Road, Wilmington, DE 19809 (US). PAYNE, Mark, S. [US/US]; 2408 Newell Drive, Wilmington, DE 19808 (US). TRIMBUR, Donald, E. [US/US]; 349 Orchard Avenue, Redwood City, CA 94601 (US). VALLE, Fernando [MX/US]; Apartment 212, 1015 Cadillac Way, Burlingame, CA 94010 (US).  <b>(74) Agent:</b> FLOYD, Linda, A.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		<b>(81) Designated States:</b> AU, BR, CA, CN, ID, IL, JP, KR, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHOD FOR THE PRODUCTION OF GLYCEROL BY RECOMBINANT ORGANISMS  <b>(57) Abstract</b>  Recombinant organisms are provided comprising genes encoding a glycerol-3-phosphate dehydrogenase and/or a glycerol-3-phosphatase activity useful for the production of glycerol from a variety of carbon substrates. The organisms further contain disruptions in the endogenous genes encoding proteins having glycerol kinase and glycerol dehydrogenase activities.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLEMETHOD FOR THE PRODUCTION OF GLYCEROL  
BY RECOMBINANT ORGANISMSFIELD OF INVENTION

5           The present invention relates to the field of molecular biology and the use of recombinant organisms for the production of glycerol and compounds derived from the glycerol biosynthetic pathway. More specifically the invention describes the construction of a recombinant cell for the production of glycerol and derived compounds from a carbon substrate, the cell containing foreign  
10 genes encoding proteins having glycerol-3-phosphate dehydrogenase (G3PDH) and glycerol-3-phosphatase (G3P phosphatase) activities where the endogenous genes encoding the glycerol-converting glycerol kinase and glycerol dehydrogenase activities have been deleted.

BACKGROUND

15           Glycerol is a compound in great demand by industry for use in cosmetics, liquid soaps, food, pharmaceuticals, lubricants, anti-freeze solutions, and in numerous other applications. The esters of glycerol are important in the fat and oil industry. Historically, glycerol has been isolated from animal fat and similar sources; however, the process is laborious and inefficient. Microbial  
20 production of glycerol is preferred.

          Not all organisms have a natural capacity to synthesize glycerol. However, the biological production of glycerol is known for some species of bacteria, algae, and yeast. The bacteria *Bacillus licheniformis* and *Lactobacillus lyopersica* synthesize glycerol. Glycerol production is found in the halotolerant  
25 algae *Dunaliella sp.* and *Asteromonas gracilis* for protection against high external salt concentrations (Ben-Amotz et al., (1982) *Experientia* 38:49-52). Similarly, various osmotolerant yeast synthesize glycerol as a protective measure. Most strains of *Saccharomyces* produce some glycerol during alcoholic fermentation and this production can be increased by the application of  
30 osmotic stress (Albertyn et al., (1994) *Mol. Cell. Biol.* 14, 4135-4144). Earlier this century glycerol was produced commercially with *Saccharomyces* cultures to which steering reagents were added such as sulfites or alkalis. Through the formation of an inactive complex, the steering agents block or inhibit the conversion of acetaldehyde to ethanol; thus, excess reducing equivalents  
35 (NADH) are available to or "steered" towards dihydroxyacetone phosphate (DHAP) for reduction to produce glycerol. This method is limited by the partial inhibition of yeast growth that is due to the sulfites. This limitation can be partially overcome by the use of alkalis which create excess NADH equivalents by a different mechanism. In this practice, the alkalis initiated a Cannizarro

disproportionation to yield ethanol and acetic acid from two equivalents of acetaldehyde. Thus, although production of glycerol is possible from naturally occurring organisms, production is often subject to the need to control osmotic stress of the cultures and the production of sulfites. A method free from these  
5 limitations is desirable. Production of glycerol from recombinant organisms containing foreign genes encoding key steps in the glycerol biosynthetic pathway is one possible route to such a method.

A number of the genes involved in the glycerol biosynthetic pathway have been isolated. For example, the gene encoding glycerol-3-phosphate  
10 dehydrogenase (DAR1, GPD1) has been cloned and sequenced from *Saccharomyces diastaticus* (Wang et al., (1994), *J. Bact.* 176:7091-7095). The DAR1 gene was cloned into a shuttle vector and used to transform *E. coli* where expression produced active enzyme. Wang et al., *supra*, recognizes that DAR1 is regulated by the cellular osmotic environment but does not suggest how the  
15 gene might be used to enhance glycerol production in a recombinant organism.

Other glycerol-3-phosphate dehydrogenase enzymes have been isolated. For example, sn-glycerol-3-phosphate dehydrogenase has been cloned and sequenced from *S. cerevisiae* (Larason et al., (1993) *Mol. Microbiol.*, 10:1101). Albertyn et al., (1994) *Mol. Cell. Biol.*, 14:4135) teach the cloning of GPD1  
20 encoding a glycerol-3-phosphate dehydrogenase from *S. cerevisiae*. Like Wang et al., both Albertyn et al. and Larason et al. recognize the osmo-sensitivity of the regulation of this gene but do not suggest how the gene might be used in the production of glycerol in a recombinant organism.

As with G3PDH, glycerol-3-phosphatase has been isolated from  
25 *Saccharomyces cerevisiae* and the protein identified as being encoded by the GPP1 and GPP2 genes (Norbeck et al., (1996) *J. Biol. Chem.*, 271:13875). Like the genes encoding G3PDH, it appears that GPP2 is osmotically-induced.

Although the genes encoding G3PDH and G3P phosphatase have been isolated, there is no teaching in the art that demonstrates glycerol production  
30 from recombinant organisms with G3PDH/G3P phosphatase expressed together or separately. Further, there is no teaching to suggest that efficient glycerol production from any wild-type organism is possible using these two enzyme activities that does not require applying some stress (salt or an osmolyte) to the cell. In fact, the art suggests that G3PDH activities may not affect glycerol  
35 production. For example, Eustace ((1987), *Can. J. Microbiol.*, 33:112-117)) teaches hybridized yeast strains that produced glycerol at greater levels than the parent strains. However, Eustace also demonstrates that G3PDH activity remained constant or slightly lower in the hybridized strains as opposed to the wild type.



Glycerol is an industrially useful material. However, other compounds may be derived from the glycerol biosynthetic pathway that also have commercial significance. For example, glycerol-producing organisms may be engineered to produce 1,3-propanediol (U.S. 5686276), a monomer having potential utility in the production of polyester fibers and the manufacture of polyurethanes and cyclic compounds. It is known for example that in some organisms, glycerol is converted to 3-hydroxypropionaldehyde and then to 1,3-propanediol through the actions of a dehydratase enzyme and an oxidoreductase enzyme, respectively. Bacterial strains able to produce 1,3-propanediol have been found, for example, in the groups *Citrobacter*, *Clostridium*, *Enterobacter*, *Ilyobacter*, *Klebsiella*, *Lactobacillus*, and *Pelobacter*. Glycerol dehydratase and diol dehydratase systems are described by Seyfried et al. (1996) *J. Bacteriol.* 178:5793-5796 and Tobimatsu et al. (1995) *J. Biol. Chem.* 270:7142-7148, respectively. Recombinant organisms, containing exogenous dehydratase enzyme, that are able to produce 1,3-propanediol have been described (U.S. 5686276). Although these organisms produce 1,3-propanediol, it is clear that they would benefit from a system that would minimize glycerol conversion.

There are a number of advantages in engineering a glycerol-producing organism for the production of 1,3-propanediol where conversion of glycerol is minimized. A microorganism capable of efficiently producing glycerol under physiological conditions is industrially desirable, especially when the glycerol itself will be used as a substrate *in vivo* as part of a more complex catabolic or biosynthetic pathway that could be perturbed by osmotic stress or the addition of steering agents (e.g., the production of 1,3-propanediol). Some attempts at creating glycerol kinase and glycerol dehydrogenase mutants have been made. For example, De Koning et al. (1990) *Appl. Microbiol Biotechnol.* 32:693-698 report the methanol-dependent production of dihydroxyacetone and glycerol by mutants of the methylotrophic yeast *Hansenula polymorpha* blocked in dihydroxyacetone kinase and glycerol kinase. Methanol and an additional substrate, required to replenish the xyulose-5-phosphate co-substrate of the assimilation reaction, were used to produce glycerol; however, a dihydroxyacetone reductase (glycerol dehydrogenase) is also required. Similarly, Shaw and Cameron, Book of Abstracts, 211th ACS National Meeting, New Orleans, LA, March 24-28 (1996), BIOT-154 Publisher: American Chemical Society, Washington, D. C., investigate the deletion of *ldhA* (lactate dehydrogenase), *glpK* (glycerol kinase), and *tpiA* (triosephosphate isomerase) for the optimization of 1,3-propanediol production. They do not suggest the expression of cloned genes for G3PDH or G3P phosphatase for the

production of glycerol or 1,3-propanediol and they do not discuss the impact of glycerol dehydrogenase.

The problem to be solved, therefore, is the lack of a process to direct carbon flux towards glycerol production by the addition or enhancement of certain enzyme activities, especially G3PDH and G3P phosphatase which respectively catalyze the conversion of dihydroxyacetone phosphate (DHAP) to glycerol-3-phosphate (G3P) and then to glycerol. The problem is complicated by the need to control the carbon flux away from glycerol by deletion or decrease of certain enzyme activities, especially glycerol kinase and glycerol dehydrogenase which respectively catalyze the conversion of glycerol plus ATP to G3P and glycerol to dihydroxyacetone (or glyceraldehyde).

#### SUMMARY OF THE INVENTION

The present invention provides a method for the production of glycerol from a recombinant organism comprising: transforming a suitable host cell with an expression cassette comprising either one or both of (a) a gene encoding a protein having glycerol-3-phosphate dehydrogenase activity and (b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity, where the suitable host cell contains a disruption in either one or both of (a) a gene encoding an endogenous glycerol kinase and (b) a gene encoding an endogenous glycerol dehydrogenase, wherein the disruption prevents the expression of active gene product; culturing the transformed host cell in the presence of at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and single-carbon substrates, whereby glycerol is produced; and recovering the glycerol produced.

The present invention further provides a process for the production of 1,3-propanediol from a recombinant organism comprising: transforming a suitable host cell with an expression cassette comprising either one or both of (a) a gene encoding a protein having glycerol-3-phosphate dehydrogenase activity and (b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity, the suitable host cell having at least one gene encoding a protein having a dehydratase activity and having a disruption in either one or both of (a) a gene encoding an endogenous glycerol kinase and (b) a gene encoding an endogenous glycerol dehydrogenase, wherein the disruption in the genes of (a) or (b) prevents the expression of active gene product; culturing the transformed host cell in the presence of at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and single-carbon substrates whereby 1,3-propanediol is produced; and recovering the 1,3-propanediol produced.

Additionally, the invention provides for a process for the production of 1,3-propanediol from a recombinant organism where multiple copies of endogeneous genes are introduced.

Further embodiments of the invention include host cells transformed with heterologous genes for the glycerol pathway as well as host cells which contain endogeneous genes for the glycerol pathway.

Additionally, the invention provides recombinant cells suitable for the production either glycerol or 1,3-propanediol, the host cells having genes expressing either one or both of a glycerol-3-phosphate dehydrogenase activity and a glycerol-3-phosphate phosphatase activity wherein the cell also has disruptions in either one or both of a gene encoding an endogenous glycerol kinase and a gene encoding an endogenous glycerol dehydrogenase, wherein the disruption in the genes prevents the expression of active gene product.

#### BRIEF DESCRIPTION OF THE FIGURES, BIOLOGICAL DEPOSITS AND SEQUENCE LISTING

Figure 1 illustrates the representative enzymatic pathways involving glycerol metabolism.

Applicants have made the following biological deposits under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for the Purposes of Patent Procedure:

Depositor Identification Reference	Int'l. Depository Designation	Date of Deposit
<i>Escherichia coli</i> pAH21/DH5 $\alpha$ (containing the GPP2 gene)	ATCC 98187	26 September 1996
<i>Escherichia coli</i> (pDAR1A/AA200) (containing the DAR1 gene)	ATCC 98248	6 November 1996
FM5 <i>Escherichia coli</i> RJF10m (containing a <i>glpK</i> disruption)	ATCC 98597	25 November 1997
FM5 <i>Escherichia coli</i> MSP33.6 (containing a <i>gldA</i> disruption)	ATCC 98598	25 November 1997

"ATCC" refers to the American Type Culture Collection international depository located at 12301 Parklawn Drive, Rockville, MD 20852 U.S.A. The designation is the accession number of the deposited material.

Applicants have provided 43 sequences in conformity with the Rules for the Standard Representation of Nucleotide and Amino Acid Sequences in Patent Applications (Annexes I and II to the Decision of the President of the EPO, published in Supplement No. 2 to OJ EPO, 12/1992) and with 37 C.F.R. 1.821-1.825 and Appendices A and B (Requirements for Application Disclosures Containing Nucleotides and/or Amino Acid Sequences).

### DETAILED DESCRIPTION OF THE INVENTION

The present invention solves the problem stated above by providing a method for the biological production of glycerol from a fermentable carbon source in a recombinant organism. The method provides a rapid, inexpensive and environmentally-responsible source of glycerol useful in the cosmetics and pharmaceutical industries. The method uses a microorganism containing cloned homologous or heterologous genes encoding glycerol-3-phosphate dehydrogenase (G3PDH) and/or glycerol-3-phosphatase (G3P phosphatase). These genes are expressed in a recombinant host having disruptions in genes encoding endogenous glycerol kinase and/or glycerol dehydrogenase enzymes. The method is useful for the production of glycerol, as well as any end products for which glycerol is an intermediate. The recombinant microorganism is contacted with a carbon source and cultured and then glycerol or any end products derived therefrom are isolated from the conditioned media. The genes may be incorporated into the host microorganism separately or together for the production of glycerol.

Applicants' process has not previously been described for a recombinant organism and required the isolation of genes encoding the two enzymes and their subsequent expression in a host cell having disruptions in the endogenous kinase and dehydrogenase genes. It will be appreciated by those familiar with this art that Applicants' process may be generally applied to the production compounds where glycerol is a key intermediate, e.g., 1,3-propanediol.

As used herein the following terms may be used for interpretation of the claims and specification.

The terms "glycerol-3-phosphate dehydrogenase" and "G3PDH" refer to a polypeptide responsible for an enzyme activity that catalyzes the conversion of dihydroxyacetone phosphate (DHAP) to glycerol-3-phosphate (G3P). *In vivo* G3PDH may be NADH; NADPH; or FAD-dependent. The NADH-dependent enzyme (EC 1.1.1.8) is encoded, for example, by several genes including GPD1 (GenBank Z74071x2), or GPD2 (GenBank Z35169x1), or GPD3 (GenBank G984182), or DAR1 (GenBank Z74071x2). The NADPH-dependent enzyme (EC 1.1.1.94) is encoded by *gpsA* (GenBank U321643, (cds 197911-196892) G466746 and L45246). The FAD-dependent enzyme (EC 1.1.99.5) is encoded by GUT2 (GenBank Z47047x23), or *glpD* (GenBank G147838), or *glpABC* (GenBank M20938).

The terms "glycerol-3-phosphatase", "sn-glycerol-3-phosphatase", or "d,l-glycerol phosphatase", and "G3P phosphatase" refer to a polypeptide responsible for an enzyme activity that catalyzes the conversion of glycerol-3-phosphate and water to glycerol and inorganic phosphate. G3P phosphatase is

encoded, for example, by GPP1 (GenBank Z47047x125), or GPP2 (GenBank U18813x11).

5 The term "glycerol kinase" refers to a polypeptide responsible for an enzyme activity that catalyzes the conversion of glycerol and ATP to glycerol-3-phosphate and ADP. The high energy phosphate donor ATP may be replaced by physiological substitutes (e.g. phosphoenolpyruvate). Glycerol kinase is encoded, for example, by GUT1 (GenBank U11583x19) and *glpK* (GenBank L19201).

10 The term "glycerol dehydrogenase" refers to a polypeptide responsible for an enzyme activity that catalyzes the conversion of glycerol to dihydroxyacetone (E.C. 1.1.1.6) or glycerol to glyceraldehyde (E.C. 1.1.1.72). A polypeptide responsible for an enzyme activity that catalyzes the conversion of glycerol to dihydroxyacetone is also referred to as a "dihydroxyacetone reductase". Glycerol dehydrogenase may be dependent upon NADH  
15 (E.C. 1.1.1.6), NADPH (E.C. 1.1.1.72), or other cofactors (e.g., E.C. 1.1.99.22). A NADH-dependent glycerol dehydrogenase is encoded, for example, by *gldA* (GenBank U00006).

The term "dehydratase enzyme" will refer to any enzyme that is capable of isomerizing or converting a glycerol molecule to the product  
20 3-hydroxypropion-aldehyde. For the purposes of the present invention the dehydratase enzymes include a glycerol dehydratase (E.C. 4.2.1.30) and a diol dehydratase (E.C. 4.2.1.28) having preferred substrates of glycerol and 1,2-propanediol, respectively. In *Citrobacter freundii*, for example, glycerol dehydratase is encoded by three polypeptides whose gene sequences are  
25 represented by *dhaB*, *dhaC* and *dhaE* (GenBank U09771: base pairs 8556-10223, 10235-10819, and 10822-11250, respectively). In *Klebsiella oxytoca*, for example, diol dehydratase is encoded by three polypeptides whose gene sequences are represented by *pddA*, *pddB*, and *pddC* (GenBank D45071: base pairs 121-1785, 1796-2470, and 2485-3006, respectively).

30 The terms "GPD1", "DAR1", "OSG1", "D2830", and "YDL022W" will be used interchangeably and refer to a gene that encodes a cytosolic glycerol-3-phosphate dehydrogenase and is characterized by the base sequence given as SEQ ID NO:1.

The term "GPD2" refers to a gene that encodes a cytosolic glycerol-3-phosphate dehydrogenase and is characterized by the base sequence given in  
35 SEQ ID NO:2.

The terms "GUT2" and "YIL155C" are used interchangeably and refer to a gene that encodes a mitochondrial glycerol-3-phosphate dehydrogenase and is characterized by the base sequence given in SEQ ID NO:3.



The terms "GPP1", "RHR2" and "YIL053W" are used interchangeably and refer to a gene that encodes a cytosolic glycerol-3-phosphatase and is characterized by the base sequence given in SEQ ID NO:4.

5 The terms "GPP2", "HOR2" and "YER062C" are used interchangeably and refer to a gene that encodes a cytosolic glycerol-3-phosphatase and is characterized by the base sequence given as SEQ ID NO:5.

The term "GUT1" refers to a gene that encodes a cytosolic glycerol kinase and is characterized by the base sequence given as SEQ ID NO:6. The term "*glpK*" refers to another gene that encodes a glycerol kinase and is  
10 characterized by the base sequence given in GeneBank L19201, base pairs 77347-78855.

The term "*gldA*" refers to a gene that encodes a glycerol dehydrogenase and is characterized by the base sequence given in GeneBank U00006, base pairs 3174-4316. The term "*dhaD*" refers to another gene that encodes a  
15 glycerol dehydrogenase and is characterized by the base sequence given in GeneBank U09771, base pairs 2557-3654.

As used herein, the terms "function" and "enzyme function" refer to the catalytic activity of an enzyme in altering the energy required to perform a specific chemical reaction. Such an activity may apply to a reaction in  
20 equilibrium where the production of both product and substrate may be accomplished under suitable conditions.

The terms "polypeptide" and "protein" are used interchangeably.

The terms "carbon substrate" and "carbon source" refer to a carbon source capable of being metabolized by host organisms of the present invention  
25 and particularly mean carbon sources selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and one-carbon substrates or mixtures thereof.

"Conversion" refers to the metabolic processes of an organism or cell that by means of a chemical reaction degrades or alters the complexity of a  
30 chemical compound or substrate.

The terms "host cell" and "host organism" refer to a microorganism capable of receiving foreign or heterologous genes and additional copies of endogeneous genes and expressing those genes to produce an active gene product.

35 The terms "production cell" and "production organism" refer to a cell engineered for the production of glycerol or compounds that may be derived from the glycerol biosynthetic pathway. The production cell will be recombinant and contain either one or both of a gene that encodes a protein having a glycerol-3-phosphate dehydrogenase activity and a gene encoding a



protein having a glycerol-3-phosphatase activity. In addition to the G3PDH and G3P phosphatase genes, the host cell will contain disruptions in one or both of a gene encoding an endogenous glycerol kinase and a gene encoding an endogenous glycerol dehydrogenase. Where the production cell is designed to produce 1,3-propanediol, it will additionally contain a gene encoding a protein having a dehydratase activity.

The terms "foreign gene", "foreign DNA", "heterologous gene", and "heterologous DNA" all refer to genetic material native to one organism that has been placed within a different host organism.

The term "endogenous" as used herein with reference to genes or polypeptides expressed by genes, refers to genes or polypeptides that are native to a production cell and are not derived from another organism. Thus an "endogenous glycerol kinase" and an "endogenous glycerol dehydrogenase" are terms referring to polypeptides encoded by genes native to the production cell.

The terms "recombinant organism" and "transformed host" refer to any organism transformed with heterologous or foreign genes. The recombinant organisms of the present invention express foreign genes encoding G3PDH and G3P phosphatase for the production of glycerol from suitable carbon substrates. Additionally, the terms "recombinant organism" and "transformed host" refer to any organism transformed with endogenous (or homologous) genes so as to increase the copy number of the genes.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding) and following (3' non-coding) the coding region. The terms "native" and "wild-type" gene refer to the gene as found in nature with its own regulatory sequences.

The terms "encoding" and "coding" refer to the process by which a gene, through the mechanisms of transcription and translation, produces an amino acid sequence. The process of encoding a specific amino acid sequence is meant to include DNA sequences that may involve base changes that do not cause a change in the encoded amino acid, or which involve base changes which may alter one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. Therefore, the invention encompasses more than the specific exemplary sequences. Modifications to the sequence, such as deletions, insertions, or substitutions in the sequence which produce silent changes that do not substantially affect the functional properties of the resulting protein molecule are also contemplated. For example, alterations in the gene sequence which reflect the degeneracy of the genetic code, or which result in the production of a chemically equivalent amino acid at a given site, are contemplated; thus, a codon for the amino acid alanine, a

hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a biologically equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the protein molecule would also not be expected to alter the activity of the protein. In some cases, it may in fact be desirable to make mutants of the sequence in order to study the effect of alteration on the biological activity of the protein. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity in the encoded products. Moreover, the skilled artisan recognizes that sequences encompassed by this invention are also defined by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65 °C), with the sequences exemplified herein.

The term "expression" refers to the transcription and translation to gene product from a gene coding for the sequence of the gene product.

The terms "plasmid", "vector", and "cassette" as used herein refer to an extra chromosomal element often carrying genes which are not part of the central metabolism of the cell and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome integrating sequences, phage or nucleotide sequences, linear or circular, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell. "Transformation cassette" refers to a specific vector containing a foreign gene and having elements in addition to the foreign gene that facilitate transformation of a particular host cell. "Expression cassette" refers to a specific vector containing a foreign gene and having elements in addition to the foreign gene that allow for enhanced expression of that gene in a foreign host.

The terms "transformation" and "transfection" refer to the acquisition of new genes in a cell after the incorporation of nucleic acid. The acquired genes may be integrated into chromosomal DNA or introduced as extrachromosomal replicating sequences. The term "transformant" refers to the cell resulting from a transformation.

The term "genetically altered" refers to the process of changing hereditary material by transformation or mutation. The terms "disruption" and "gene interrupt" as applied to genes refer to a method of genetically altering an organism by adding to or deleting from a gene a significant portion of that gene such that the protein encoded by that gene is either not expressed or not expressed in active form.

#### Glycerol Biosynthetic Pathway

It is contemplated that glycerol may be produced in recombinant organisms by the manipulation of the glycerol biosynthetic pathway found in most microorganisms. Typically, a carbon substrate such as glucose is converted to glucose-6-phosphate via hexokinase in the presence of ATP. Glucose-phosphate isomerase catalyzes the conversion of glucose-6-phosphate to fructose-6-phosphate and then to fructose-1,6-diphosphate through the action of 6-phosphofructokinase. The diphosphate is then taken to dihydroxyacetone phosphate (DHAP) via aldolase. Finally NADH-dependent G3PDH converts DHAP to glycerol-3-phosphate which is then dephosphorylated to glycerol by G3P phosphatase. (Agarwal (1990), *Adv. Biochem. Engrg.* 41:114).

#### Genes encoding G3PDH, glycerol dehydrogenase, G3P phosphatase and glycerol kinase

The present invention provides genes suitable for the expression of G3PDH and G3P phosphatase activities in a host cell.

Genes encoding G3PDH are known. For example, GPD1 has been isolated from *Saccharomyces* and has the base sequence given by SEQ ID NO:1, encoding the amino acid sequence given in SEQ ID NO:7 (Wang et al., *supra*). Similarly, G3PDH activity has also been isolated from *Saccharomyces* encoded by GPD2 having the base sequence given in SEQ ID NO:2 encoding the amino acid sequence given in SEQ ID NO:8 (Eriksson et al., (1995) *Mol. Microbiol.*, 17:95).

For the purposes of the present invention it is contemplated that any gene encoding a polypeptide responsible for G3PDH activity is suitable wherein that activity is capable of catalyzing the conversion of dihydroxyacetone phosphate (DHAP) to glycerol-3-phosphate (G3P). Further, it is contemplated that any gene encoding the amino acid sequence of G3PDH as given by SEQ ID NOS:7, 8, 9, 10, 11 and 12 corresponding to the genes GPD1, GPD2, GUT2, gpaA, glpD, and the  $\alpha$  subunit of glpABC respectively, will be functional in the present invention wherein that amino acid sequence may encompass amino acid substitutions, deletions or additions that do not alter the function of the enzyme. The skilled person will appreciate that genes encoding G3PDH isolated from other sources will also be suitable for use in the present invention. For

example, genes isolated from prokaryotes include GenBank accessions M34393, M20938, L06231, U12567, L45246, L45323, L45324, L45325, U32164, U32689, and U39682. Genes isolated from fungi include GenBank accessions U30625, U30676 and X56162; genes isolated from insects include GenBank  
 5 accessions X61223 and X14179; and genes isolated from mammalian sources include GenBank accessions U12424, M25558 and X78593.

Genes encoding G3P phosphatase are known. For example, GPP2 has been isolated from *Saccharomyces cerevisiae* and has the base sequence given by SEQ ID NO:5, which encodes the amino acid sequence given in SEQ ID NO:13  
 10 (Norbeck et al., (1996), *J. Biol. Chem.*, 271:13875).

For the purposes of the present invention, any gene encoding a G3P phosphatase activity is suitable for use in the method wherein that activity is capable of catalyzing the conversion of glycerol-3-phosphate and water to glycerol and inorganic phosphate. Further, any gene encoding the amino acid  
 15 sequence of G3P phosphatase as given by SEQ ID NOS:13 and 14 corresponding to the genes GPP2 and GPP1 respectively, will be functional in the present invention including any amino acid sequence that encompasses amino acid substitutions, deletions or additions that do not alter the function of the G3P phosphatase enzyme. The skilled person will appreciate that genes encoding  
 20 G3P phosphatase isolated from other sources will also be suitable for use in the present invention. For example, the dephosphorylation of glycerol-3-phosphate to yield glycerol may be achieved with one or more of the following general or specific phosphatases: alkaline phosphatase (EC 3.1.3.1) [GenBank M19159, M29663, U02550 or M33965]; acid phosphatase (EC 3.1.3.2) [GenBank  
 25 U51210, U19789, U28658 or L20566]; glycerol-3-phosphatase (EC 3.1.3.-) [GenBank Z38060 or U18813x11]; glucose-1-phosphatase (EC 3.1.3.10) [GenBank M33807]; glucose-6-phosphatase (EC 3.1.3.9) [GenBank U00445]; fructose-1,6-bisphosphatase (EC 3.1.3.11) [GenBank X12545 or J03207] or phosphatidyl glycerol phosphate phosphatase (EC 3.1.3.27) [GenBank M23546  
 30 and M23628].

Genes encoding glycerol kinase are known. For example, GUT1 encoding the glycerol kinase from *Saccharomyces* has been isolated and sequenced (Pavlik et al. (1993), *Curr. Genet.*, 24:21) and the base sequence is given by SEQ ID NO:6, which encodes the amino acid sequence given in  
 35 SEQ ID NO:15. Alternatively, *glpK* encodes a glycerol kinase from *E. coli* and is characterized by the base sequence given in GeneBank L19201, base pairs 77347-78855.

Genes encoding glycerol dehydrogenase are known. For example, *gldA* encodes a glycerol dehydrogenase from *E. coli* and is characterized by the base

sequence given in GeneBank U00006, base pairs 3174-4316. Alternatively, *dhaD* refers to another gene that encodes a glycerol dehydrogenase from *Citrobacter freundii* and is characterized by the base sequence given in GeneBank U09771, base pairs 2557-3654.

5 Host cells

Suitable host cells for the recombinant production of glycerol by the expression of G3PDH and G3P phosphatase may be either prokaryotic or eukaryotic and will be limited only by their ability to express active enzymes. Preferred host cells will be those bacteria, yeasts, and filamentous fungi typically useful for the production of glycerol such as *Citrobacter*, *Enterobacter*, *Clostridium*, *Klebsiella*, *Aerobacter*, *Lactobacillus*, *Aspergillus*, *Saccharomyces*, *Schizosaccharomyces*, *Zygosaccharomyces*, *Pichia*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Torulopsis*, *Methylobacter*, *Escherichia*, *Salmonella*, *Bacillus*, *Streptomyces* and *Pseudomonas*. Preferred in the present invention are *E. coli* and *Saccharomyces*.

Where glycerol is a key intermediate in the production of 1,3-propanediol the host cell will either have an endogenous gene encoding a protein having a dehydratase activity or will acquire such a gene through transformation. Host cells particularly suited for production of 1,3-propanediol are *Citrobacter*, *Enterobacter*, *Clostridium*, *Klebsiella*, *Aerobacter*, *Lactobacillus*, and *Salmonella*, which have endogenous genes encoding dehydratase enzymes. Additionally, host cells that lack such an endogeneous gene include *E. coli*.

20 Vectors And Expression Cassettes

The present invention provides a variety of vectors and transformation and expression cassettes suitable for the cloning, transformation and expression of G3PDH and G3P phosphatase into a suitable host cell. Suitable vectors will be those which are compatible with the bacterium employed. Suitable vectors can be derived, for example, from a bacteria, a virus (such as bacteriophage T7 or a M-13 derived phage), a cosmid, a yeast or a plant. Protocols for obtaining and using such vectors are known to those in the art (Sambrook et al., Molecular Cloning: A Laboratory Manual - volumes 1, 2, 3 (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY, 1989)).

Typically, the vector or cassette contains sequences directing transcription and translation of the appropriate gene, a selectable marker, and sequences allowing autonomous replication or chromosomal integration. Suitable vectors comprise a region 5' of the gene which harbors transcriptional initiation controls and a region 3' of the DNA fragment which controls transcriptional termination. It is most preferred when both control regions are derived from genes homologous to the transformed host cell. Such control



regions need not be derived from the genes native to the specific species chosen as a production host.

Initiation control regions, or promoters, which are useful to drive expression of the G3PDH and G3P phosphatase genes in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving these genes is suitable for the present invention including but not limited to CYC1, HIS3, GAL1, GAL10, ADH1, PGK, PHO5, GAPDH, ADC1, TRP1, URA3, LEU2, ENO, and TPI (useful for expression in *Saccharomyces*); AOX1 (useful for expression in *Pichia*); and lac, trp,  $\lambda P_L$ ,  $\lambda P_R$ , T7, tac, and trc, (useful for expression in *E. coli*).

Termination control regions may also be derived from various genes native to the preferred hosts. Optionally, a termination site may be unnecessary; however, it is most preferred if included.

For effective expression of the instant enzymes, DNA encoding the enzymes are linked operably through initiation codons to selected expression control regions such that expression results in the formation of the appropriate messenger RNA.

#### Transformation Of Suitable Hosts And Expression Of G3PDH And G3P Phosphatase For The Production Of Glycerol

Once suitable cassettes are constructed they are used to transform appropriate host cells. Introduction of the cassette containing the genes encoding G3PDH and/or G3P phosphatase into the host cell may be accomplished by known procedures such as by transformation, e.g., using calcium-permeabilized cells, electroporation, or by transfection using a recombinant phage virus (Sambrook et al., *supra*).

In the present invention AH21 and DAR1 cassettes were used to transform the *E. coli* DH5 $\alpha$  and FM5 as fully described in the GENERAL METHODS and EXAMPLES.

Alternatively, it is contemplated that suitable host cells comprising endogenous G3PDH and/or G3P phosphatase genes may be manipulated so that the relevant genes are upregulated for the production of glycerol.

Methods for upregulation of endogenous genes are well known in the art. For example, to upregulate the desired gene(s), a structural gene is generally placed downstream from a promoter region on the DNA which is recognized by the recipient microorganism. In addition to the promoter, one may include other regulatory sequences that increase or control expression from heterologous genes. In addition, one may alter the regulatory sequences of endogenous genes by any known genetic manipulation for the same purpose. Expression may be controlled



by an inducer or a repressor so that the microorganism coordinately expresses the gene(s) necessary to complete the desired metabolic pathway.

In the instant invention host cells containing endogenous genes encoding G3PDH and/or G3P phosphatase activities could be placed under the control of regulated promoters (e.g. *lac* or *osmy*) or constitutive promoters. For example, a cassette may be constructed to contain a specific inducible or constitutive promoter, flanked by DNA of sufficient length and homology to the native gene to permit targeting. Introduction of the cassette under suitable growth conditions will result in homologous recombination between the cassette and the targeted portion of the gene and the replacement of the relevant native promoter with the regulatable promoter. Such methods may be employed to effect the upregulation of endogenous genes encoding G3PDH and/or G3P phosphatase activities for the production of glycerol.

Random And Site Specific Mutagenesis For Disrupting Enzyme Activities:

Enzyme pathways by which organisms metabolize glycerol are known in the art, Figure 1. Glycerol is converted to glycerol-3-phosphate (G3P) by an ATP-dependent glycerol kinase; the G3P may then be oxidized to DHAP by G3PDH. In a second pathway, glycerol is oxidized to dihydroxyacetone (DHA) by a glycerol dehydrogenase; the DHA may then be converted to DHAP by an ATP-dependent DHA kinase. In a third pathway, glycerol is oxidized to glyceraldehyde by a glycerol dehydrogenase; the glyceraldehyde may be phosphorylated to glyceraldehyde-3-phosphate by an ATP-dependent kinase. DHAP and glyceraldehyde-3-phosphate, interconverted by the action of triosephosphate isomerase, may be further metabolized via central metabolism pathways. These pathways, by introducing by-products, are deleterious to glycerol production.

One aspect of the present invention is the ability to provide a production organism for the production of glycerol where the glycerol-converting activities of glycerol kinase and glycerol dehydrogenase have been deleted. Methods of creating deletion mutants are common and well known in the art. For example, wild type cells may be exposed to a variety of agents such as radiation or chemical mutagens and then screened for the desired phenotype. When creating mutations through radiation either ultraviolet (UV) or ionizing radiation may be used. Suitable short wave UV wavelengths for genetic mutations will fall within the range of 200 nm to 300 nm where 254 nm is preferred. UV radiation in this wavelength principally causes changes within nucleic acid sequence from guanine and cytosine to adenine and thymidine. Since all cells have DNA repair mechanisms that would repair most UV induced mutations, agents such as caffeine and other inhibitors may be added to interrupt the repair process and

maximize the number of effective mutations. Long wave UV mutations using light in the 300 nm to 400 nm range are also possible but are generally not as effective as the short wave UV light unless used in conjunction with various activators such as psoralen dyes that interact with the DNA.

5 Mutagenesis with chemical agents is also effective for generating mutants and commonly used substances include chemicals that affect nonreplicating DNA such as  $\text{HNO}_2$  and  $\text{NH}_2\text{OH}$ , as well as agents that affect replicating DNA such as acridine dyes, notable for causing frameshift mutations. Specific methods for creating mutants using radiation or chemical agents are well  
10 documented in the art. See for example Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition (1989) Sinauer Associates, Inc., Sunderland, MA., or Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36, 227, (1992), herein incorporated by reference.

After mutagenesis has occurred, mutants having the desired phenotype  
15 may be selected by a variety of methods. Random screening is most common where the mutagenized cells are selected for the ability to produce the desired product or intermediate. Alternatively, selective isolation of mutants can be performed by growing a mutagenized population on selective media where only resistant colonies can develop. Methods of mutant selection are highly  
20 developed and well known in the art of industrial microbiology. See Brock, *Supra.*, DeMancilha et al., *Food Chem.*, 14, 313, (1984).

Biological mutagenic agents which target genes randomly are well known in the art. See for example De Bruijn and Rossbach in *Methods for General and Molecular Bacteriology* (1994) American Society for Microbiology,  
25 Washington, D.C. Alternatively, provided that gene sequence is known, chromosomal gene disruption with specific deletion or replacement is achieved by homologous recombination with an appropriate plasmid. See for example Hamilton et al. (1989) *J. Bacteriol.* 171:4617-4622, Balbas et al. (1993) *Gene* 136:211-213, Gueldener et al. (1996) *Nucleic Acids Res.* 24:2519-2524, and  
30 Smith et al. (1996) *Methods Mol. Cell. Biol.* 5:270-277.

It is contemplated that any of the above cited methods may be used for the deletion or inactivation of glycerol kinase and glycerol dehydrogenase activities in the preferred production organism.

#### Media and Carbon Substrates

35 Fermentation media in the present invention must contain suitable carbon substrates. Suitable substrates may include but are not limited to mono-saccharides such as glucose and fructose, oligosaccharides such as lactose or sucrose, polysaccharides such as starch or cellulose or mixtures thereof and unpurified mixtures from renewable feedstocks such as cheese whey permeate,

cornsteep liquor, sugar beet molasses, and barley malt. Additionally, the carbon substrate may also be one-carbon substrates such as carbon dioxide, or methanol for which metabolic conversion into key biochemical intermediates has been demonstrated.

5 Glycerol production from single carbon sources (e.g., methanol, formaldehyde or formate) has been reported in methylotrophic yeasts (Yamada et al. (1989), *Agric. Biol. Chem.*, 53(2):541-543) and in bacteria (Hunter et al. (1985), *Biochemistry*, 24:4148-4155). These organisms can assimilate single carbon compounds, ranging in oxidation state from methane to formate, and  
10 produce glycerol. The pathway of carbon assimilation can be through ribulose monophosphate, through serine, or through xylulose-monophosphate (Gottschalk, Bacterial Metabolism, Second Edition, Springer-Verlag: New York (1986)). The ribulose monophosphate pathway involves the condensation of formate with ribulose-5-phosphate to form a 6 carbon sugar that becomes  
15 fructose and eventually the three carbon product, glyceraldehyde-3-phosphate. Likewise, the serine pathway assimilates the one-carbon compound into the glycolytic pathway via methylenetetrahydrofolate.

In addition to one and two carbon substrates, methylotrophic organisms are also known to utilize a number of other carbon-containing compounds such  
20 as methylamine, glucosamine and a variety of amino acids for metabolic activity. For example, methylotrophic yeast are known to utilize the carbon from methylamine to form trehalose or glycerol (Bellion et al. (1993), *Microb. Growth C1 Compd.*, [Int. Symp.], 7th, 415-32. Editor(s): Murrell, J. Collin; Kelly, Don P. Publisher: Intercept, Andover, UK). Similarly, various species  
25 of *Candida* will metabolize alanine or oleic acid (Sulter et al. (1990), *Arch. Microbiol.*, 153(5):485-9). Hence, the source of carbon utilized in the present invention may encompass a wide variety of carbon-containing substrates and will only be limited by the choice of organism.

Although all of the above mentioned carbon substrates and mixtures  
30 thereof are suitable in the present invention, preferred carbon substrates are monosaccharides, oligosaccharides, polysaccharides, single-carbon substrates or mixtures thereof. More preferred are sugars such as glucose, fructose, sucrose, maltose, lactose and single carbon substrates such as methanol and carbon dioxide. Most preferred as a carbon substrate is glucose.

35 In addition to an appropriate carbon source, fermentation media must contain suitable minerals, salts, cofactors, buffers and other components, known to those skilled in the art, suitable for the growth of the cultures and promotion of the enzymatic pathway necessary for glycerol production.

### Culture Conditions

Typically cells are grown at 30 °C in appropriate media. Preferred growth media are common commercially prepared media such as Luria Bertani (LB) broth, Sabouraud Dextrose (SD) broth, or Yeast medium (YM) broth.

- 5 Other defined or synthetic growth media may also be used and the appropriate medium for growth of the particular microorganism will be known by one skilled in the art of microbiology or fermentation science. The use of agents known to modulate catabolite repression directly or indirectly, e.g., cyclic adenosine 3':5'-monophosphate, may also be incorporated into the reaction
- 10 media. Similarly, the use of agents known to modulate enzymatic activities (e.g., sulfites, bisulfites, and alkalis) that lead to enhancement of glycerol production may be used in conjunction with or as an alternative to genetic manipulations.

- Suitable pH ranges for the fermentation are between pH 5.0 to pH 9.0 where the range of pH 6.0 to pH 8.0 is preferred for the initial condition.
- 15

Reactions may be performed under aerobic or anaerobic conditions where anaerobic or microaerobic conditions are preferred.

### Identification of G3PDH, glycerol dehydrogenase, G3P phosphatase, and glycerol kinase activities

- 20 The levels of expression of the proteins G3PDH, G3P phosphatase glycerol dehydrogenase, and glycerol kinase are measured by enzyme assays. Generally, G3PDH activity and glycerol dehydrogenase activity assays rely on the spectral properties of the cosubstrate, NADH, in the DHAP conversion to G-3-P and the DHA conversion to glycerol, respectively. NADH has intrinsic
- 25 UV/vis absorption and its consumption can be monitored spectrophotometrically at 340 nm. G3P phosphatase activity can be measured by any method of measuring the inorganic phosphate liberated in the reaction. The most commonly used detection method uses the visible spectroscopic determination of a blue-colored phosphomolybdate ammonium complex. Glycerol kinase activity
- 30 can be measured by the detection of G3P from glycerol and ATP, for example, by NMR. Assays can be directed toward more specific characteristics of individual enzymes if necessary, for example, by the use of alternate cofactors.

### Identification and recovery of glycerol and other products (e.g. 1,3-propanediol)

- Glycerol and other products (e.g. 1,3-propanediol) may be identified and
- 35 quantified by high performance liquid chromatography (HPLC) and gas chromatography/mass spectroscopy (GC/MS) analyses on the cell-free extracts. Preferred is a HPLC method where the fermentation media are analyzed on an analytical ion exchange column using a mobile phase of 0.01N sulfuric acid in an isocratic fashion.

Methods for the recovery of glycerol from fermentation media are known in the art. For example, glycerol can be obtained from cell media by subjecting the reaction mixture to the following sequence of steps: filtration; water removal; organic solvent extraction; and fractional distillation (U.S. Patent  
5 No. 2,986,495).

#### Description Of The Preferred Embodiments

##### Production of Glycerol

The present invention describes a method for the production of glycerol from a suitable carbon source utilizing a recombinant organism. Particularly  
10 suitable in the invention is a bacterial host cell, transformed with an expression cassette carrying either or both of a gene that encodes a protein having a glycerol-3-phosphate dehydrogenase activity and a gene encoding a protein having a glycerol-3-phosphatase activity. In addition to the G3PDH and G3P phosphatase genes, the host cell will contain disruptions in either or both of  
15 genes encoding endogenous glycerol kinase and glycerol dehydrogenase enzymes. The combined effect of the foreign G3PDH and G3P phosphatase genes (providing a pathway from the carbon source to glycerol) with the gene disruptions (blocking the conversion of glycerol) results in an organism that is capable of efficient and reliable glycerol production.

20 Although the optimal organism for glycerol production contains the above mentioned gene disruptions, glycerol production is possible with a host cell containing either one or both of the foreign G3PDH and G3P phosphatase genes in the absence of such disruptions. For example, the recombinant *E. coli* strain AA200 carrying the DAR1 gene (Example 1) was capable of producing  
25 between 0.38 g/L and 0.48 g/L of glycerol depending on fermentation parameters. Similarly, the *E. coli* DH5 $\alpha$ , carrying and expressible GPP2 gene (Example 2), was capable of 0.2 g/L of glycerol production. Where both genes are present, (Example 3 and 4), glycerol production attained about 40 g/L. Where both genes are present in conjunction with an elimination of the  
30 endogenous glycerol kinase activity, a reduction in the conversion of glycerol may be seen (Example 8). Furthermore, the presence of glycerol dehydrogenase activity is linked to the conversion of glycerol under glucose-limited conditions; thus, it is anticipated that the elimination of glycerol dehydrogenase activity will result in the reduction of glycerol conversion (Example 8).

##### 35 Production of 1,3-propanediol

The present invention may also be adapted for the production of 1,3-propanediol by utilizing recombinant organisms expressing the foreign G3PDH and/or G3P phosphatase genes and containing disruptions in the endogenous glycerol kinase and/or glycerol dehydrogenase activities.



Additionally, the invention provides for the process for the production of 1,3-propanediol from a recombinant organism where multiple copies of endogeneous genes are introduced. In addition to these genetic alterations, the production cell will require the presence of a gene encoding an active  
5 dehydratase enzyme. The dehydratase enzyme activity may either be a glycerol dehydratase or a diol dehydratase. The dehydratase enzyme activity may result from either the expression of an endogenous gene or from the expression of a foreign gene transfected into the host organism. Isolation and expression of genes encoding suitable dehydratase enzymes are well known in the art and are  
10 taught by applicants in PCT/US96/06705, filed 5 November 1996 and U.S. 5686276 and U.S. 5633362, hereby incorporated by reference. It will be appreciated that, as glycerol is a key intermediate in the production of 1,3-propanediol, where the host cell contains a dehydratase activity in conjunction with expressed foreign G3PDH and/or G3P phosphatase genes and  
15 in the absence of the glycerol-converting glycerol kinase or glycerol dehydrogenase activities, the cell will be particularly suited for the production of 1,3-propanediol.

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred  
20 embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

25

### EXAMPLES

#### GENERAL METHODS

Procedures for phosphorylations, ligations, and transformations are well known in the art. Techniques suitable for use in the following examples may be found in Sambrook et al., Molecular Cloning: A Laboratory Manual, Second  
30 Edition, Cold Spring Harbor Laboratory Press (1989).

Materials and methods suitable for the maintenance and growth of bacterial cultures are well known in the art. Techniques suitable for use in the following examples may be found in Manual of Methods for General Bacteriology (Phillipp Gerhardt, R. G. E. Murray, Ralph N. Costilow, Eugene  
35 W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, eds), American Society for Microbiology, Washington, DC. (1994) or in Biotechnology: A Textbook of Industrial Microbiology (Thomas D. Brock, Second Edition (1989) Sinauer Associates, Inc., Sunderland, MA). All reagents and materials used for the growth and maintenance of bacterial cells were



obtained from Aldrich Chemicals (Milwaukee, WI), DIFCO Laboratories (Detroit, MI), GIBCO/BRL (Gaithersburg, MD), or Sigma Chemical Company (St. Louis, MO) unless otherwise specified.

The meaning of abbreviations is as follows: "h" means hour(s), "min" means minute(s), "sec" means second(s), "d" means day(s), "mL" means milliliters, "L" means liters.

#### Cell strains

The following *Escherichia coli* strains were used for transformation and expression of G3PDH and G3P phosphatase. Strains were obtained from the *E. coli* Genetic Stock Center, ATCC, or Life Technologies (Gaithersburg, MD).

AA200 (*garB10 fhuA22 ompF627 fadL701 relA1 pit-10 spoT1 tpi-1 phoM510 mcrB1*) (Anderson et al., (1970), *J. Gen. Microbiol.*, 62:329).

BB20 (*tonA22 ΔphoA8 fadL701 relA1 glpR2 glpD3 pit-10 gpsA20 spoT1 T2R*) (Cronan et al., *J. Bact.*, 118:598).

DH5α (*deoR endA1 gyrA96 hsdR17 recA1 relA1 supE44 thi-1 Δ(lacZYA-argFV169) phi80lacZΔM15 F<sup>-</sup>*) (Woodcock et al., (1989), *Nucl. Acids Res.*, 17:3469).

FM5 *Escherichia coli* (ATCC 53911)

#### Identification of Glycerol

The conversion of glucose to glycerol was monitored by HPLC and/or GC. Analyses were performed using standard techniques and materials available to one of skill in the art of chromatography. One suitable method utilized a Waters Maxima 820 HPLC system using UV (210 nm) and RI detection. Samples were injected onto a Shodex SH-1011 column (8 mm x 300 mm; Waters, Milford, MA) equipped with a Shodex SH-1011P precolumn (6 mm x 50 mm), temperature-controlled at 50 °C, using 0.01 N H<sub>2</sub>SO<sub>4</sub> as mobile phase at a flow rate of 0.69 mL/min. When quantitative analysis was desired, samples were prepared with a known amount of trimethylacetic acid as an external standard. Typically, the retention times of 1,3-propanediol (RI detection), glycerol (RI detection) and glucose (RI detection) were 21.39 min, 17.03 min and 12.66 min, respectively.

Glycerol was also analyzed by GC/MS. Gas chromatography with mass spectrometry detection for separation and quantitation of glycerol was performed using a DB-WAX column (30 m, 0.32 mm I.D., 0.25 μm film thickness, J & W Scientific, Folsom, CA) at the following conditions: injector: split, 1:15;

sample volume: 1  $\mu$ L; temperature profile: 150 °C initial temperature with 30 sec hold, 40 °C/min to 180 °C, 20 °C/min to 240 °C, hold for 2.5 min. Detection: EI Mass Spectrometry (Hewlett Packard 5971, San Fernando, CA), quantitative SIM using ions 61 m/z and 64 m/z as target ions for glycerol and glycerol-d8, and ion 43 m/z as qualifier ion for glycerol. Glycerol-d8 was used as an internal standard.

#### Assay for glycerol-3-phosphatase, G3P phosphatase

The assay for enzyme activity was performed by incubating the extract with an organic phosphate substrate in a bis-Tris or MES and magnesium buffer, pH 6.5. The substrate used was either l- $\alpha$ -glycerol phosphate, or d,l- $\alpha$ -glycerol phosphate. The final concentrations of the reagents in the assay are: buffer (20 mM bis-Tris or 50 mM MES);  $MgCl_2$  (10 mM); and substrate (20 mM). If the total protein in the sample was low and no visible precipitation occurs with an acid quench, the sample was conveniently assayed in the cuvette. This method involved incubating an enzyme sample in a cuvette that contained 20 mM substrate (50  $\mu$ L, 200 mM), 50 mM MES, 10 mM  $MgCl_2$ , pH 6.5 buffer. The final phosphatase assay volume was 0.5 mL. The enzyme-containing sample was added to the reaction mixture; the contents of the cuvette were mixed and then the cuvette was placed in a circulating water bath at  $T = 37$  °C for 5 to 120 min, the length of time depending on whether the phosphatase activity in the enzyme sample ranged from 2 to 0.02 U/mL. The enzymatic reaction was quenched by the addition of the acid molybdate reagent (0.4 mL). After the Fiske SubbaRow reagent (0.1 mL) and distilled water (1.5 mL) were added, the solution was mixed and allowed to develop. After 10 min, to allow full color development, the absorbance of the samples was read at 660 nm using a Cary 219 UV/Vis spectrophotometer. The amount of inorganic phosphate released was compared to a standard curve that was prepared by using a stock inorganic phosphate solution (0.65 mM) and preparing 6 standards with final inorganic phosphate concentrations ranging from 0.026 to 0.130  $\mu$ mol/mL.

#### Spectrophotometric Assay for Glycerol 3-Phosphate Dehydrogenase (G3PDH) Activity

The following procedure was used as modified below from a method published by Bell et al. (1975), *J. Biol. Chem.*, 250:7153-8. This method involved incubating an enzyme sample in a cuvette that contained 0.2 mM NADH; 2.0 mM dihydroxyacetone phosphate (DHAP), and enzyme in 0.1 M Tris/HCl, pH 7.5 buffer with 5 mM DTT, in a total volume of 1.0 mL at 30 °C. The spectrophotometer was set to monitor absorbance changes at the fixed wavelength of 340 nm. The instrument was blanked on a cuvette containing

buffer only. After the enzyme was added to the cuvette, an absorbance reading was taken. The first substrate, NADH (50  $\mu$ L 4 mM NADH; absorbance should increase approx 1.25 AU), was added to determine the background rate. The rate should be followed for at least 3 min. The second substrate, DHAP (50  $\mu$ L 40 mM DHAP), was then added and the absorbance change over time was monitored for at least 3 min to determine the gross rate. G3PDH activity was defined by subtracting the background rate from the gross rate.

#### $^{13}$ C-NMR Assay for Glycerol Kinase Activity

An appropriate amount of enzyme, typically a cell-free crude extract, was added to a reaction mixture containing 40 mM ATP, 20 mM  $\text{MgSO}_4$ , 21 mM uniformly  $^{13}\text{C}$  labelled glycerol (99%, Cambridge Isotope Laboratories), and 0.1 M Tris-HCl, pH 9 for 75 min at 25  $^{\circ}\text{C}$ . The conversion of glycerol to glycerol 3-phosphate was detected by  $^{13}\text{C}$ -NMR (125 MHz): glycerol (63.11 ppm, d,  $J = 41$  Hz and 72.66 ppm, t,  $J = 41$  Hz); glycerol 3-phosphate (62.93 ppm, d,  $J = 41$  Hz; 65.31 ppm, br d,  $J = 43$  Hz; and 72.66 ppm, dt,  $J = 6, 41$  Hz).

#### NADH-linked Glycerol Dehydrogenase Assay

NADH -linked glycerol dehydrogenase activity in *E. coli* strains (*gldA*) was determined after protein separation by non-denaturing polyacrylamide gel electrophoresis. The conversion of glycerol plus  $\text{NAD}^+$  to dihydroxyacetone plus NADH was coupled with the conversion of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) to a deeply colored formazan, using phenazine methosulfate (PMS) as mediator. (Tang et al. (1997) *J. Bacteriol.* 140:182).

Electrophoresis was performed in duplicate by standard procedures using native gels (8-16% TG, 1.5 mm, 15 lane gels from Novex, San Diego, CA). Residual glycerol was removed from the gels by washing 3x with 50 mM Tris or potassium carbonate buffer, pH 9 for 10 min. The duplicate gels were developed, with and without glycerol (approx. 0.16 M final concentration), in 15 mL of assay solution containing 50 mM Tris or potassium carbonate, pH 9, 60 mg ammonium sulfate, 75 mg  $\text{NAD}^+$ , 1.5 mg MTT, and 0.5 mg PMS.

The presence or absence of NADH -linked glycerol dehydrogenase activity in *E. coli* strains (*gldA*) was also determined, following polyacrylamide gel electrophoresis, by reaction with polyclonal antibodies raised to purified *K. pneumoniae* glycerol dehydrogenase (*dhaD*).

## PLASMID CONSTRUCTION AND STRAIN CONSTRUCTION

### Cloning and expression of glycerol 3-phosphatase for increase of glycerol production in *E. coli* DH5 $\alpha$ and FM5

The *Saccharomyces cerevisiae* chromosome V lamda clone 6592 (Gene Bank, accession # U18813x11) was obtained from ATCC. The glycerol 3-phosphate phosphatase (GPP2) gene was cloned by cloning from the lamda clone as target DNA using synthetic primers (SEQ ID NO:16 with SEQ ID NO:17) incorporating an BamHI-RBS-XbaI site at the 5' end and a SmaI site at the 3' end. The product was subcloned into pCR-Script (Stratagene, Madison, WI) at the SrfI site to generate the plasmids pAH15 containing GPP2. The plasmid pAH15 contains the GPP2 gene in the inactive orientation for expression from the lac promoter in pCR-Script SK+. The BamHI-SmaI fragment from pAH15 containing the GPP2 gene was inserted into pBlueScriptII SK+ to generate plasmid pAH19. The pAH19 contains the GPP2 gene in the correct orientation for expression from the lac promoter. The XbaI-PstI fragment from pAH19 containing the GPP2 gene was inserted into pPHOX2 to create plasmid pAH21. The pAH21/ DH5 $\alpha$  is the expression plasmid.

### Plasmids for the over-expression of DAR1 in *E. coli*

DAR1 was isolated by PCR cloning from genomic *S. cerevisiae* DNA using synthetic primers (SEQ ID NO:18 with SEQ ID NO:19). Successful PCR cloning places an NcoI site at the 5' end of DAR1 where the ATG within NcoI is the DAR1 initiator methionine. At the 3' end of DAR1 a BamHI site is introduced following the translation terminator. The PCR fragments were digested with NcoI + BamHI and cloned into the same sites within the expression plasmid pTrc99A (Pharmacia, Piscataway, NJ) to give pDAR1A.

In order to create a better ribosome binding site at the 5' end of DAR1, an SpeI-RBS-NcoI linker obtained by annealing synthetic primers (SEQ ID NO:20 with SEQ ID NO:21) was inserted into the NcoI site of pDAR1A to create pAH40. Plasmid pAH40 contains the new RBS and DAR1 gene in the correct orientation for expression from the trc promoter of pTrc99A (Pharmacia, Piscataway, NJ). The NcoI-BamHI fragment from pDAR1A and an second set of SpeI-RBS-NcoI linker obtained by annealing synthetic primers (SEQ ID NO:22 with SEQ ID NO:23) was inserted into the SpeI-BamHI site of pBC-SK+ (Stratagene, Madison, WI) to create plasmid pAH42. The plasmid pAH42 contains a chloramphenicol resistant gene.

### Construction of expression cassettes for DAR1 and GPP2

Expression cassettes for DAR1 and GPP2 were assembled from the individual DAR1 and GPP2 subclones described above using standard molecular biology methods. The BamHI-PstI fragment from pAH19 containing the

ribosomal binding site (RBS) and GPP2 gene was inserted into pAH40 to create pAH43. The BamHI-PstI fragment from pAH19 containing the RBS and GPP2 gene was inserted into pAH42 to create pAH45.

The ribosome binding site at the 5' end of GPP2 was modified as follows. A BamHI-RBS-SpeI linker, obtained by annealing synthetic primers GATCCAGGAAACAGA (SEQ ID NO:24) with CTAGTCTGTTTCCTG (SEQ ID NO:25) to the XbaI-PstI fragment from pAH19 containing the GPP2 gene, was inserted into the BamHI-PstI site of pAH40 to create pAH48. Plasmid pAH48 contains the DAR1 gene, the modified RBS, and the GPP2 gene in the correct orientation for expression from the trc promoter of pTrc99A (Pharmacia, Piscataway, NJ).

#### Transformation of *E. coli*

All the plasmids described here were transformed into *E. coli* DH5 $\alpha$  or FM5 using standard molecular biology techniques. The transformants were verified by its DNA RFLP pattern.

### EXAMPLE 1

#### PRODUCTION OF GLYCEROL FROM *E. COLI* TRANSFORMED WITH G3PDH GENE

##### Media

Synthetic media was used for anaerobic or aerobic production of glycerol using *E. coli* cells transformed with pDAR1A. The media contained per liter 6.0 g Na<sub>2</sub>HPO<sub>4</sub>, 3.0 g KH<sub>2</sub>PO<sub>4</sub>, 1.0 g NH<sub>4</sub>Cl, 0.5 g NaCl, 1 mL 20% MgSO<sub>4</sub>·7H<sub>2</sub>O, 8.0 g glucose, 40 mg casamino acids, 0.5 ml 1% thiamine hydrochloride, 100 mg ampicillin.

##### Growth Conditions

Strain AA200 harboring pDAR1A or the pTrc99A vector was grown in aerobic conditions in 50 mL of media shaking at 250 rpm in 250 mL flasks at 37 °C. At A<sub>600</sub> 0.2-0.3 isopropylthio- $\beta$ -D-galactoside was added to a final concentration of 1 mM and incubation continued for 48 h. For anaerobic growth samples of induced cells were used to fill Falcon #2054 tubes which were capped and gently mixed by rotation at 37 °C for 48 h. Glycerol production was determined by HPLC analysis of the culture supernatants. Strain pDAR1A/AA200 produced 0.38 g/L glycerol after 48 h under anaerobic conditions, and 0.48 g/L under aerobic conditions.



EXAMPLE 2  
PRODUCTION OF GLYCEROL FROM *E. COLI*  
TRANSFORMED WITH G3P PHOSPHATASE GENE (GPP2)

Media

5 Synthetic phoA media was used in shake flasks to demonstrate the increase of glycerol by GPP2 expression in *E. coli*. The phoA medium contained per liter: Amisoy, 12 g; ammonium sulfate, 0.62 g; MOPS, 10.5 g; Na-citrate, 1.2 g; NaOH (1 M), 10 mL; 1 M MgSO<sub>4</sub>, 12 mL; 100X trace elements, 12 mL; 50% glucose, 10 mL; 1% thiamine, 10 mL; 100 mg/mL  
10 L-proline, 10 mL; 2.5 mM FeCl<sub>3</sub>, 5 mL; mixed phosphates buffer, 2 mL (5 mL 0.2 M NaH<sub>2</sub>PO<sub>4</sub> + 9 mL 0.2 M K<sub>2</sub>HPO<sub>4</sub>), and pH to 7.0. The 100X traces elements for phoA medium /L contained: ZnSO<sub>4</sub>·7 H<sub>2</sub>O, 0.58 g; MnSO<sub>4</sub>·H<sub>2</sub>O, 0.34 g; CuSO<sub>4</sub>·5 H<sub>2</sub>O, 0.49 g; CoCl<sub>2</sub>·6 H<sub>2</sub>O, 0.47 g; H<sub>3</sub>BO<sub>3</sub>, 0.12 g, NaMoO<sub>4</sub>·2 H<sub>2</sub>O, 0.48 g.

15 Shake Flasks Experiments

The strains pAH21/DH5α (containing GPP2 gene) and pPHOX2/DH5α (control) were grown in 45 mL of media (phoA media, 50 ug/mL carbenicillin, and 1 ug/mL vitamin B<sub>12</sub>) in a 250 mL shake flask at 37 °C. The cultures were grown under aerobic condition (250 rpm shaking) for 24 h. Glycerol production  
20 was determined by HPLC analysis of the culture supernatant. pAH21/DH5α produced 0.2 g/L glycerol after 24 h.

EXAMPLE 3  
PRODUCTION OF GLYCEROL FROM D-GLUCOSE USING  
RECOMBINANT *E. COLI* CONTAINING BOTH GPP2 AND DAR1

25 Growth for demonstration of increased glycerol production by *E. coli* DH5α-containing pAH43 proceeds aerobically at 37 °C in shake-flask cultures (erlenmeyer flasks, liquid volume 1/5th of total volume).

Cultures in minimal media/1 % glucose shake-flasks are started by inoculation from overnight LB/1 % glucose culture with antibiotic selection.  
30 Minimal media are: filter-sterilized defined media, final pH 6.8 (HCl), contained per liter: 12.6 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 13.7 g K<sub>2</sub>HPO<sub>4</sub>, 0.2 g yeast extract (Difco), 1 g NaHCO<sub>3</sub>, 5 mg vitamin B<sub>12</sub>, 5 mL Modified Balch's Trace-Element Solution (the composition of which can be found in Methods for General and  
35 Molecular Bacteriology (P. Gerhardt et al., eds, p. 158, American Society for Microbiology, Washington, DC (1994)). The shake-flasks are incubated at 37 °C with vigorous shaking for overnight, after which they are sampled for GC analysis of the supernatant. The pAH43/DH5α showed glycerol production of 3.8 g/L after 24 h.



EXAMPLE 4PRODUCTION OF GLYCEROL FROM D-GLUCOSE USING  
RECOMBINANT *E. COLI* CONTAINING BOTH GPP2 AND DAR1

Example 4 illustrates the production of glucose from the recombinant

5 *E. coli* DH5 $\alpha$ /pAH48, containing both the GPP2 and DAR1 genes.

The strain DH5 $\alpha$ /pAH48 was constructed as described above in the  
GENERAL METHODS.

Pre-Culture

DH5 $\alpha$ /pAH48 were pre-cultured for seeding into a fermentation run.

10 Components and protocols for the pre-culture are listed below.

Pre-Culture Media

	KH <sub>2</sub> PO <sub>4</sub>	30.0 g/L
	Citric acid	2.0 g/L
	MgSO <sub>4</sub> ·7H <sub>2</sub> O	2.0 g/L
15	98% H <sub>2</sub> SO <sub>4</sub>	2.0 mL/L
	Ferric ammonium citrate	0.3 g/L
	CaCl <sub>2</sub> ·2H <sub>2</sub> O	0.2 g/L
	Yeast extract	5.0 g/L
	Trace metals	5.0 mL/L
20	Glucose	10.0 g/L
	Carbenicillin	100.0 mg/L

The above media components were mixed together and the pH adjusted  
to 6.8 with NH<sub>4</sub>OH. The media was then filter sterilized.

Trace metals were used according to the following recipe:

25	Citric acid, monohydrate	4.0 g/L
	MgSO <sub>4</sub> ·7H <sub>2</sub> O	3.0 g/L
	MnSO <sub>4</sub> ·H <sub>2</sub> O	0.5 g/L
	NaCl	1.0 g/L
	FeSO <sub>4</sub> ·7H <sub>2</sub> O	0.1 g/L
30	CoCl <sub>2</sub> ·6H <sub>2</sub> O	0.1 g/L
	CaCl <sub>2</sub>	0.1 g/L
	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.1 g/L
	CuSO <sub>4</sub> ·5 H <sub>2</sub> O	10 mg/L
	AlK(SO <sub>4</sub> ) <sub>2</sub> ·12H <sub>2</sub> O	10 mg/L
35	H <sub>3</sub> BO <sub>3</sub>	10 mg/L
	Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	10 mg/L
	NiSO <sub>4</sub> ·6H <sub>2</sub> O	10 mg/L
	Na <sub>2</sub> SeO <sub>3</sub>	10 mg/L
	Na <sub>2</sub> WO <sub>4</sub> ·2H <sub>2</sub> O	10 mg/L

Cultures were started from seed culture inoculated from 50  $\mu$ L frozen stock (15% glycerol as cryoprotectant) to 600 mL medium in a 2-L Erlenmeyer flask. Cultures were grown at 30 °C in a shaker at 250 rpm for approximately 12 h and then used to seed the fermenter.

5 Fermentation growth

Vessel

15-L stirred tank fermenter

Medium

	$\text{KH}_2\text{PO}_4$	6.8 g/L
10	Citric acid	2.0 g/L
	$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	2.0 g/L
	98% $\text{H}_2\text{SO}_4$	2.0 mL/L
	Ferric ammonium citrate	0.3 g/L
	$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	0.2 g/L
15	Mazu DF204 antifoam	1.0 mL/L

The above components were sterilized together in the fermenter vessel. The pH was raised to 6.7 with  $\text{NH}_4\text{OH}$ . Yeast extract (5 g/L) and trace metals solution (5 mL/L) were added aseptically from filter sterilized stock solutions. Glucose was added from 60% feed to give final concentration of 10 g/L.

20 Carbenicillin was added at 100 mg/L. Volume after inoculation was 6 L.

Environmental Conditions For Fermentation

The temperature was controlled at 36 °C and the air flow rate was controlled at 6 standard liters per minute. Back pressure was controlled at 0.5 bar. The agitator was set at 350 rpm. Aqueous ammonia was used to  
25 control pH at 6.7. The glucose feed (60% glucose monohydrate) rate was controlled to maintain excess glucose.

Results

The results of the fermentation run are given in Table 1.

Table 1

EFT (hr)	OD550 (AU)	[Glucose] (g/L)	[Glycerol] (g/L)	Total Glucose Fed (g)	Total Glycerol Produced (g)
0	0.8	9.3		25	
6	4.7	4.0	2.0	49	14
8	5.4	0	3.6	71	25
10	6.7	0.0	4.7	116	33
12	7.4	2.1	7.0	157	49
14.2	10.4	0.3	10.0	230	70
16.2	18.1	9.7	15.5	259	106
18.2	12.4	14.5		305	
20.2	11.8	17.4	17.7	353	119
22.2	11.0	12.6		382	
24.2	10.8	6.5	26.6	404	178
26.2	10.9	6.8		442	
28.2	10.4	10.3	31.5	463	216
30.2	10.2	13.1	30.4	493	213
32.2	10.1	8.1	28.2	512	196
34.2	10.2	3.5	33.4	530	223
36.2	10.1	5.8		548	
38.2	9.8	5.1	36.1	512	233

5

EXAMPLE 5

ENGINEERING OF GLYCEROL KINASE MUTANTS OF *E. COLI* FM5  
FOR PRODUCTION OF GLYCEROL FROM GLUCOSE

Construction of integration plasmid for glycerol kinase gene replacement in  
*E. coli* FM5

- 10 *E. coli* FM5 genomic DNA was prepared using the Puregene DNA Isolation Kit (Gentra Systems, Minneapolis, MN). A 1.0 kb DNA fragment containing partial *glpF* and glycerol kinase (*glpK*) genes was amplified by PCR (Mullis and Faloona, *Methods Enzymol.*, 155:335-350, 1987) from FM5 genomic DNA using primers SEQ ID NO:26 and SEQ ID NO:27. A 1.1 kb
- 15 DNA fragment containing partial *glpK* and *glpX* genes was amplified by PCR from FM5 genomic DNA using primers SEQ ID NO:28 and SEQ ID NO:29. A *MunI* site was incorporated into primer SEQ ID NO:28. The 5' end of primer SEQ ID NO:28 was the reverse complement of primer SEQ ID NO:27 to enable

subsequent overlap extension PCR. The gene splicing by overlap extension technique (Horton et al., *BioTechniques*, 8:528-535, 1990) was used to generate a 2.1 kb fragment by PCR using the above two PCR fragments as templates and primers SEQ ID NO:26 and SEQ ID NO:29. This fragment represented a deletion of 0.8 kb from the central region of the 1.5 kb *glpK* gene. Overall, this fragment had 1.0 kb and 1.1 kb flanking regions on either side of the *MunI* cloning site (within the partial *glpK*) to allow for chromosomal gene replacement by homologous recombination.

The above 2.1 kb PCR fragment was blunt-ended (using mung bean nuclease) and cloned into the pCR-Blunt vector using the Zero Blunt PCR Cloning Kit (Invitrogen, San Diego, CA) to yield the 5.6 kb plasmid pRN100 containing kanamycin and Zeocin resistance genes. The 1.2 kb *HincII* fragment from pLoxCat1 (unpublished results), containing a chloramphenicol-resistance gene flanked by bacteriophage P1 *loxP* sites (Snaith et al., *Gene*, 166:173-174, 1995), was used to interrupt the *glpK* fragment in plasmid pRN100 by ligating it to *MunI*-digested (and blunt-ended) plasmid pRN100 to yield the 6.9 kb plasmid pRN101-1. A 376 bp fragment containing the R6K origin was amplified by PCR from the vector pGP704 (Miller and Mekalanos, *J. Bacteriol.*, 170:2575-2583, 1988) using primers SEQ ID NO:30 and SEQ ID NO:31, blunt-ended, and ligated to the 5.3 kb *Asp718-AatII* fragment (which was blunt-ended) from pRN101-1 to yield the 5.7 kb plasmid pRN102-1 containing kanamycin and chloramphenicol resistance genes. Substitution of the *ColE1* origin region in pRN101-1 with the R6K origin to generate pRN102-1 also involved deletion of most of the Zeocin resistance gene. The host for pRN102-1 replication was *E. coli* SY327 (Miller and Mekalanos, *J. Bacteriol.*, 170:2575-2583, 1988) which contains the *pir* gene necessary for the function of the R6K origin.

#### Engineering Of Glycerol Kinase Mutant RJF10m With Chloramphenicol Resistance Gene Interrupt

*E. coli* FM5 was electrotransformed with the non-replicative integration plasmid pRN102-1 and transformants that were chloramphenicol-resistant (12.5 µg/mL) and kanamycin-sensitive (30 µg/mL) were further screened for glycerol non-utilization on M9 minimal medium containing 1 mM glycerol. An *EcoRI* digest of genomic DNA from one such mutant, RJF10m, when probed with the intact *glpK* gene via Southern analysis (Southern, *J. Mol. Biol.*, 98:503-517, 1975) indicated that it was a double-crossover integrant (*glpK* gene replacement) since the two expected 7.9 kb and 2.0 kb bands were observed, owing to the presence of an additional *EcoRI* site within the chloramphenicol resistance gene. The wild-type control yielded the single expected 9.4 kb band.

A  $^{13}\text{C}$  NMR analysis of mutant RJF10m confirmed that it was incapable of converting  $^{13}\text{C}$ -labeled glycerol and ATP to glycerol-3-phosphate. This *glpK* mutant was further analyzed by genomic PCR using primer combinations SEQ ID NO:32 and SEQ ID NO:33, SEQ ID NO:34 and SEQ ID NO:35, and SEQ ID NO:32 and SEQ ID NO:35 which yielded the expected 2.3 kb, 2.4 kb, and 4.0 kb PCR fragments respectively. The wild-type control yielded the expected 3.5 kb band with primers SEQ ID NO:32 and SEQ ID NO:35. The *glpK* mutant RJF10m was electrotransformed with plasmid pAH48 to allow glycerol production from glucose. The *glpK* mutant *E. coli* RJF10m has been deposited with ATCC under the terms of the Budapest Treaty on 24 November, 1997.

Engineering Of Glycerol Kinase Mutant RJF10 With Chloramphenicol Resistance Gene Interrupt Removed

After overnight growth on YENB medium (0.75% yeast extract, 0.8% nutrient broth) at 37 °C, *E. coli* RJF10m in a water suspension was electrotransformed with plasmid pJW168 (unpublished results), which contained the bacteriophage P1 Cre recombinase gene under the control of the IPTG-inducible *lacUV5* promoter, a temperature-sensitive pSC101 replicon, and an ampicillin resistance gene. Upon outgrowth in SOC medium at 30 °C, transformants were selected at 30 °C (permissive temperature for pJW168 replication) on LB agar medium supplemented with carbenicillin (50 µg/mL) and IPTG (1 mM). Two serial overnight transfers of pooled colonies were carried out at 30 °C on fresh LB agar medium supplemented with carbenicillin and IPTG in order to allow excision of the chromosomal chloramphenicol resistance gene via recombination at the *loxP* sites mediated by the Cre recombinase (Hoess and Abremski, *J. Mol. Biol.*, 181:351-362, 1985). Resultant colonies were replica-plated on to LB agar medium supplemented with carbenicillin and IPTG and LB agar supplemented with chloramphenicol (12.5 µg/mL) to identify colonies that were carbenicillin-resistant and chloramphenicol-sensitive indicating marker gene removal. An overnight 30 °C culture of one such colony was used to inoculate 10 mL of LB medium. Upon growth at 30 °C to OD (600 nm) of 0.6, the culture was incubated at 37 °C overnight. Several dilutions were plated on prewarmed LB agar medium and the plates incubated overnight at 42 °C (the non-permissive temperature for pJW168 replication). Resultant colonies were replica-plated on to LB agar medium and LB agar medium supplemented with carbenicillin (75 µg/mL) to identify colonies that were carbenicillin-sensitive indicating loss of plasmid pJW168. One such *glpK* mutant, RJF10, was further analyzed by genomic PCR using primers SEQ ID NO:32 and SEQ ID NO:35 and yielded the expected 3.0 kb band confirming marker gene excision. Glycerol non-utilization by mutant RJF10 was confirmed



by lack of growth on M9 minimal medium containing 1 mM glycerol. The *glpK* mutant RJF10 was electrotransformed with plasmid pAH48 to allow glycerol production from glucose.

#### EXAMPLE 6

##### 5 CONSTRUCTION OF *E. COLI* STRAIN WITH *GLDA* GENE KNOCKOUT

The *gldA* gene was isolated from *E. coli* by PCR (K. B. Mullis and F. A. Faloon (1987) Meth. Enzymol. 155:335-350) using primers SEQ ID NO:36 and SEQ ID NO:37, which incorporate terminal Sph1 and Xba1 sites, respectively, and cloned (T. Maniatis 1982 Molecular Cloning. A Laboratory  
10 Manual. Cold Spring Harbor, Cold Spring Harbor, NY) between the Sph1 and Xba1 sites in pUC18, to generate pKP8. pKP8 was cut at the unique Sal1 and Nco1 sites within the *gldA* gene, the ends flushed with Klenow and religated, resulting in a 109 bp deletion in the middle of *gldA* and regeneration of a unique Sal1 site, to generate pKP9. A 1.4 kb DNA fragment containing the gene  
15 conferring kanamycin resistance (kan), and including about 400 bps of DNA upstream of the translational start codon and about 100 bps of DNA downstream of the translational stop codon, was isolated from pET-28a(+) (Novagen, Madison, Wis) by PCR using primers SEQ ID NO:38 and SEQ ID NO:39, which incorporate terminal Sal1 sites, and subcloned into the unique Sal1 site of  
20 pKP9, to generate pKP13. A 2.1 kb DNA fragment beginning 204 bps downstream of the *gldA* translational start codon and ending 178 bps upstream of the *gldA* translational stop codon, and containing the kan insertion, was isolated from pKP13 by PCR using primers SEQ ID NO:40 and SEQ ID NO:41, which incorporate terminal Sph1 and Xba1 sites, respectively, was subcloned between  
25 the Sph1 and Xba1 sites in pMAK705 (Genencor International, Palo Alto, Calif.), to generate pMP33. *E. coli* FM5 was transformed with pMP33 and selected on 20 ug/mL kan at 30 °C, which is the permissive temperature for pMAK705 replication. One colony was expanded overnight at 30 °C in liquid media supplemented with 20 ug/mL kan. Approximately 32,000 cells were  
30 plated on 20 ug/mL kan and incubated for 16 hrs at 44 °C, which is the restrictive temperature for pMAK705 replication. Transformants growing at 44 °C have plasmid integrated into the chromosome, occurring at a frequency of approximately 0.0001. PCR and Southern blot (E.M. Southern 1975 *J. Mol. Biol.* 98:503-517) analyses were used to determine the nature of the  
35 chromosomal integration events in the transformants. Western blot analysis (H. Towbin, et al. (1979) *Proc. Natl. Acad. Sci.* 76:4350) was used to determine whether glycerol dehydrogenase protein, the product of *gldA*, is produced in the transformants. An activity assay was used to determine whether glycerol dehydrogenase activity remained in the transformants. Activity in

glycerol dehydrogenase bands on native gels was determined by coupling the conversion of glycerol + NAD(+) → dihydroxyacetone + NADH to the conversion of a tetrazolium dye, MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] to a deeply colored formazan, with phenazine methosulfate as mediator. Glycerol dehydrogenase also requires the presence of 30 mM ammonium sulfate and 100 mM Tris, pH 9 (C.-T. Tang, et al. (1997) *J. Bacteriol.* 140:182). Of 8 transformants analyzed, 6 were determined to be *gldA* knockouts. *E. coli* MSP33.6 has been deposited with ATCC under the terms of the Budapest Treaty on 24 November 1997.

10

EXAMPLE 7CONSTRUCTION OF *E. COLI* STRAINWITH *GLPK* AND *GLDA* GENE KNOCKOUTS

A 1.6 kb DNA fragment containing the *gldA* gene and including 228 bps of DNA upstream of the translational start codon and 220 bps of DNA downstream of the translational stop codon was isolated from *E. coli* by PCR using primers SEQ ID NO:42 and SEQ ID NO:43, which incorporate terminal SphI and XbaI sites, respectively, and cloned between the SphI and XbaI sites of pUC18, to generate pQN2. pQN2 was cut at the unique SalI and NcoI sites within the *gldA* gene, the ends flushed with Klenow and religated, resulting in a 109 bps deletion in the middle of *gldA* and regeneration of a unique SalI site, to generate pQN4. A 1.2 kb DNA fragment containing the gene conferring kanamycin resistance (kan), and flanked by loxP sites was isolated from pLoxKan2 (Genencor International, Palo Alto, Calif.) as a StuI/XhoI fragment, the ends flushed with Klenow, and subcloned into pQN4 at the SalI site after flushing with Klenow, to generate pQN8. A 0.4 kb DNA fragment containing the R6K origin of replication was isolated from pGP704 (Miller and Mekalanos, *J. Bacteriol.*, 170:2575-2583, 1988) by PCR using primers SEQ ID NO:44 and SEQ ID NO:45, which incorporate terminal SphI and XbaI sites, respectively, and ligated to the 2.8 kb SphI/XbaI DNA fragment containing the *gldA::kan* cassette from pQN8, to generate pKP22. A 1.0 kb DNA fragment containing the gene conferring chloramphenicol resistance (cam), and flanked by loxP sites was isolated from pLoxCat2 (Genencor International, Palo Alto, Calif.) as an XbaI fragment, and subcloned into pKP22 at the XbaI site, to generate pKP23. *E. coli* strain RJF10 (see EXAMPLE 5), which is *glpK*<sup>-</sup>, was transformed with pKP23 and transformants with the phenotype kanRcamS were isolated, indicating double crossover integration, which was confirmed by southern blot analysis. Glycerol dehydrogenase gel activity assays (as described in EXAMPLE 6) demonstrated that active glycerol dehydrogenase was not present in these transformants. The kan marker was removed from the chromosome using the Cre-producing plasmid

pJW168, as described in EXAMPLE 5, to produce strain KLP23. Several isolates with the phenotype kan<sup>S</sup> demonstrated no glycerol dehydrogenase activity, and southern blot analysis confirmed loss of the kan marker.

5 SEQ ID NO:44:  
CACGCATGCAGTTCAACCTGTTGATAGTAC

SEQ ID NO:45:  
GCGTCTAGATCCTTTTAAATTAAAAATG

10

#### EXAMPLE 8

#### CONSUMPTION OF GLYCEROL PRODUCED FROM D-GLUCOSE BY RECOMBINANT *E. COLI* CONTAINING BOTH GPP2 AND DAR1 WITH AND WITHOUT GLYCEROL KINASE (GLPK) ACTIVITY

EXAMPLE 8 illustrates the consumption of glycerol by the recombinant  
15 *E. coli* FM5/pAH48 and RJF10/pAH48. The strains FM5/pAH48 and  
RJF10/pAH48 were constructed as described above in the GENERAL  
METHODS.

#### Pre-Culture

FM5/pAH48 and RJF10/pAH48 were pre-cultured for seeding a  
20 fermenter in the same medium used for fermentation, or in LB supplemented  
with 1 % glucose. Either carbenicillin or ampicillin were used (100 mg/L) for  
plasmid maintenance. The medium for fermentation is as described in  
EXAMPLE 4.

Cultures were started from frozen stocks (15% glycerol as  
25 cryoprotectant) in 600 mL medium in a 2-L Erlenmeyer flask, grown at 30 °C  
in a shaker at 250 rpm for approximately 12 h, and used to seed the fermenter.

#### Fermentation growth

A 15-L stirred tank fermenter with 5-7 L initial volume was prepared as  
described in EXAMPLE 4. Either carbenicillin or ampicillin were used  
30 (100 mg/L) for plasmid maintenance.

#### Environmental Conditions to Evaluate Glycerol Kinase (GlpK) Activity

The temperature was controlled at 30 °C and the air flow rate controlled  
at 6 standard liters per minute. Back pressure was controlled at 0.5 bar.  
Dissolved oxygen tension was controlled at 10% by stirring. Aqueous ammonia  
35 was used to control pH at 6.7. The glucose feed (60% glucose) rate was  
controlled to maintain excess glucose until glycerol had accumulated to at least  
25 g/L. Glucose was then depleted, resulting in the net metabolism of glycerol.  
Table 2 shows the resulting conversion of glycerol.

Table 2  
Conversion of glycerol by FM5/pAH48 (wt) and RJF10/pAH48 (*glpK*)

Strain	number of examples	rate of glycerol consumption g/OD/hr
FM5/pAH48	2	0.095 ± 0.015
RJF10/pAH48	3	0.021 ± 0.011

As is seen by the data in Table 2, the rate of glycerol consumption decreases about 4-5 fold where endogenous glycerol kinase activity is eliminated.

5 Environmental Conditions to Evaluate Glycerol Dehydrogenase (GldA) Activity

The temperature was controlled at 30 °C and the air flow rate controlled at 6 standard liters per minute. Back pressure was controlled at 0.5 bar. Dissolved oxygen tension was controlled at 10% by stirring. Aqueous ammonia was used to control pH at 6.7. In the first fermentation, glucose was kept in excess for the duration of the fermentation. The second fermentation was operated with no residual glucose after the first 25 hours. Samples over time from the two fermentations were taken for evaluation of GlpK and GldA activities. Table 3 summarizes RJF10/pAH48 fermentations that show the effects of GldA on selectivity for glycerol.

15

Table 3  
GldA and GlpK activities from two RJF10/pAH48 fermentations

Fermentation	Time (hrs)	GldA	GlpK	Overall selectivity (g/g)
1	25	-	-	42%
	46	-	-	49%
	61	+	-	54%
2	25	+	-	41%
	46	++	-	14%
	61	++	-	12%

As is seen by the data in Table 3, the presence of glycerol dehydrogenase (GldA) activity is linked to the conversion of glycerol under glucose-limited conditions; thus, it is anticipated that eliminating glycerol dehydrogenase activity will reduce glycerol conversion.

20

EXAMPLE 9  
PRODUCTION OF GLYCEROL FROM D-GLUCOSE USING  
RECOMBINANT *E. BLATTAE* CONTAINING  
BOTH GPP2 AND DAR1

5           Example 9 illustrates the production of glycerol from D-glucose from recombinant *E. blattae* containing both GPP2 and DAR1 genes.

*E. blattae*, obtained from the ATCC and having ATCC accession number 33429, was grown at 30 °C until the culture reached an OD of about 0.6 AU at 600 nm. The culture was then transformed with pAH48, a plasmid comprising  
10   GPP2 and DAR1 genes (described in WO 98/21341), using electroporation techniques. The transformants were confirmed by DNA RFLP pattern and antibiotic resistance (200 ug/mL carbenicillin).

          The transformed *E. blattae* was grown aerobically at 35 °C in shake-flask cultures. The cultures were grown in a defined medium plus 2 % glucose  
15   with antibiotic selection and were started by inoculation from an overnight culture grown in LB plus 1 % glucose with antibiotic selection. The defined medium contained per liter: 27.2 g KH<sub>2</sub>PO<sub>4</sub>, 2 g citric acid, 2 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 1.2 ml 98 % H<sub>2</sub>SO<sub>4</sub>, 0.3 g ferric ammonium citrate, 0.2 g CaCl<sub>2</sub>·2H<sub>2</sub>O, 10 g yeast extract (Difco), 5 mL Modified Balch's Trace-Element Solution (the  
20   composition of which can be found in Methods for General and Molecular Bacteriology (P. Gerhardt et al., eds, p. 158, American Society for Microbiology, Washington, DC, (1994)). The defined medium was filter-sterilized and adjusted to a final pH 6.8 with NH<sub>4</sub>OH. The shake-flasks were incubated at 35 °C overnight with vigorous shaking. The supernatant was then  
25   subjected to HPLC analysis for the presence of glycerol. After the overnight incubation, the *E. blattae* containing pAH48 produced 7.63 g/L of glycerol. The control, which was wild-type *E. blattae* (ATCC 33429) grown under the same conditions, produced = 0.2 g/L of glycerol.

EXAMPLE 10  
PRODUCTION OF GLYCEROL FROM D-GLUCOSE  
USING RECOMBINANT *E. COLI* DEFICIENT IN *GLDA* AND *GLPK*  
AND CONTAINING BOTH GPP2 AND DAR1 INTEGRATED  
INTO THE CHROMOSOME

30           This Example illustrates the production of glycerol from D glucose from recombinant *E. coli* with *gldA* and *glpK* gene knockouts and containing both  
35   GPP2 and DAR1 encoding genes integrated into the host cell chromosome.

*E. coli* strain KLP23, prepared as described in Example 7, is deficient in both glycerol kinase (product of *glpK*) and glycerol dehydrogenase (product of



*gldA*) activities. KLP23 containing DAR1, GPP2 and a *loxP* flanked chloramphenicol resistant gene integrated into the chromosome at the *ampC* location was prepared and is referred to as AH76RIcm.

Integration plasmids were designed and constructed based on a cre-lox  
5 integration system (Hoess, supra). In order to create the integration plasmids, a Hind III - SmaI fragment of *pLoxCat1* was inserted into Hind III and Sma I linearized pAH48 to create pAH48cm2. The pAH48 plasmid contains DAR1 and GPP2 genes expressed under the control of the *trc* promoter. The 3.5 kb  
10 ApaL I fragment of pAH48cm2 was blunt ended with T4 DNA polymerase (Boehringer Mannheim Biochemical) and dNTPs and inserted into NruI linearized *pInt-ampC* (Genencor International, CA), using *E. coli* SY327 (Miller et al., *J. Bacteriol.* 170:2575-2583, 1998) as a host to create pAH76 and pAH76R. The "R" means reverse orientation of the integration cassette. Both  
15 plasmids, pAH76 and pAH76R, contain a R6K origin of replication and are not able to replicate in KLP23. The plasmids pAH76 and pAH76R were used to transform KLP23 for integration at the *ampC* location of the *E. coli* chromosome. The transformants were selected on 10 ug/ml of chloramphenicol and were kanamycin sensitive, yielding double crossover integration. These  
20 *E. coli* transformants are named AH76Icm and AH76RIcm.

AH76RIcm cultures were grown in shake-flasks in defined medium  
(described in Example 9) plus 2.5% glucose started by inoculation from an overnight LB culture having 1% glucose and antibiotic selection. The shake-  
25 flasks (erlenmeyer flasks, liquid volume 1/5<sup>th</sup> of total volume) were incubated at 37 °C with vigorous shaking overnight, after which the supernatant was sampled for glycerol using a colormetric enzyme assay (Sigma, Procedure No. 337) on a Monarch 2000 instrument (Instrumentation Laboratory Co., Lexington, MA ).  
AH77RIcm showed glycerol production of 6.7 g/L after 25 hr.

*E. coli* pAH76RI has the chloramphenicol gene deleted from AH76RIcm. The chloramphenicol gene was deleted from the chromosome using the  
30 Cre-producing plasmid, pJW168, as described in Example 5. The transformants were selected for carbenicillin resistance and chloramphenicol sensitivity under 1 mM IPTG induction at 30 °C. After removal of the chloramphenicol gene, AH76RI was grown on LB medium without any antibiotics to cure pJW168. The final version of AH76RI is not able to grow on chloramphenicol or  
35 carbenicillin selection.

AH76RI cultures were grown in shake-flasks in a defined media plus 2% glucose started by inoculation from an overnight LB/1% glucose culture. The shake-flasks were incubated at 35 °C with vigorous shaking overnight, after which the supernatant was sampled for glycerol using a colormetric assay

(Sigma, Procedure No. 337) on a Monarch 2000 instrument (Instrumentation Laboratory Co. Lexington, MA ). AH77RI showed glycerol production of 4.6 g/L after 24 hr.

- 5 All the plasmids described in this example were transformed into *E. coli* KLP23 using standard molecular biology techniques. The transformants were verified by DNA RFLP pattern, antibiotic resistance, PCR amplification, or G3P phosphatase assay.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>20</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, Virginia 20110-2209 USA	
Date of deposit 26 September 1996	Accession Number ATCC98187
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
In respect of those designations in which a European patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC)	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="text-align: right; font-weight: bold; font-size: small;">For receiving Office use only</div> <div style="padding: 5px;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border-top: 1px solid black; padding: 5px;">Authorized officer </div>	<div style="text-align: right; font-weight: bold; font-size: small;">For International Bureau use only</div> <div style="padding: 5px;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border-top: 1px solid black; padding: 5px;">Authorized officer</div>
--	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>21</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, Virginia 20110-2209 USA	
Date of deposit 6 November 1996	Accession Number ATCC98248
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
In respect of those designations in which a European patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC)	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="text-align: right; font-weight: bold; font-size: small;">For receiving Office use only</div> <div style="padding: 5px;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border-top: 1px solid black; padding: 5px;">Authorized officer </div>	<div style="text-align: right; font-weight: bold; font-size: small;">For International Bureau use only</div> <div style="padding: 5px;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border-top: 1px solid black; padding: 5px;">Authorized officer</div>
--	---

Form PCT/RO/134 (July 1992)

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>22</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, Virginia 20110-2209 USA	
Date of deposit 25 November 1997	Accession Number ATCC98597
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
In respect of those designations in which a European patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC)	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


<div style="text-align: right; font-weight: bold; font-size: small;">For receiving Office use only</div> <div style="padding: 5px;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border-top: 1px solid black; padding: 5px;">Authorized officer </div>	<div style="text-align: right; font-weight: bold; font-size: small;">For International Bureau use only</div> <div style="padding: 5px;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border-top: 1px solid black; padding: 5px;">Authorized officer</div>
--	---



## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>23</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, Virginia 20110-2209 USA	
Date of deposit 25 November 1997	Accession Number ATCC98598
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
In respect of those designations in which a European patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC)	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="border-bottom: 1px solid black; margin-bottom: 5px;">For receiving Office use only</div> <div style="display: flex; align-items: center; margin-bottom: 5px;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border-bottom: 1px solid black; margin-bottom: 5px;">Authorized officer</div> <div style="text-align: center;"></div>	<div style="border-bottom: 1px solid black; margin-bottom: 5px;">For International Bureau use only</div> <div style="display: flex; align-items: center; margin-bottom: 5px;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border-bottom: 1px solid black; margin-bottom: 5px;">Authorized officer</div>
---	--

Form PCT/RO/134 (July 1992)

## WHAT IS CLAIMED IS:

1. A method for the production of glycerol from a recombinant organism comprising:
  - (i) transforming a suitable host cell with an expression cassette comprising either one or both of
    - (a) a gene encoding a protein having glycerol-3-phosphate dehydrogenase activity, and
    - (b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity,
  - 10 the suitable host cell having a disruption in either one or both of
    - (a) an endogenous gene encoding a polypeptide having glycerol kinase activity, and
    - (b) an endogenous gene encoding a polypeptide having glycerol dehydrogenase activity,
  - 15 wherein the disruption prevents the expression of active gene product;
    - (ii) culturing the transformed host cell of (i) in the presence of at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and single-carbon substrates, whereby glycerol is produced; and
    - 20 (iii) optionally recovering the glycerol produced in (ii).
2. The method of Claim 1 wherein the expression cassette comprises a gene encoding a glycerol-3-phosphate dehydrogenase enzyme.
3. The method of Claim 1 wherein the expression cassette comprises a gene encoding a glycerol-3-phosphate phosphatase enzyme.
- 25 4. The method of Claim 1 wherein the expression cassette comprises genes encoding a glycerol-3-phosphate phosphatase enzyme and a glycerol-3-phosphate dehydrogenase enzyme.
5. The method of Claim 1 wherein the host cell contains a disruption in a gene encoding an endogenous glycerol kinase enzyme wherein the disruption prevents the expression of active gene product.
- 30 6. The method of Claim 1 wherein the host cell contains a disruption in a gene encoding an endogenous glycerol dehydrogenase enzyme wherein the disruption prevents the expression of active gene product.
7. The method of Claim 1 wherein the host cell contains a) a disruption in a gene encoding an endogenous glycerol kinase enzyme and b) a disruption in a gene encoding an endogenous glycerol dehydrogenase enzyme, wherein the disruptions in the respective genes prevent the expression of active gene product from either gene.
- 35

8. The method of Claim 1 wherein the suitable host cell is selected from the group consisting of bacteria, yeast, and filamentous fungi.

9. The method of Claim 8 wherein the suitable host cell is selected from the group consisting of *Citrobacter*, *Enterobacter*, *Clostridium*, *Klebsiella*,  
5 *Aerobacter*, *Lactobacillus*, *Aspergillus*, *Saccharomyces*, *Schizosaccharomyces*,  
*Zygosaccharomyces*, *Pichia*, *Kluyveromyces*, *Candida*, *Hansenula*,  
*Debaryomyces*, *Mucor*, *Torulopsis*, *Methylobacter*, *Escherichia*, *Salmonella*,  
*Bacillus*, *Streptomyces*, and *Pseudomonas*.

10. The method of Claim 9 wherein the suitable host cell is *E. coli* or  
10 *Saccharomyces sp.*

11. The method of Claim 1 wherein the carbon source is glucose.

12. The method of Claim 1 wherein the protein having glycerol-3-phosphate dehydrogenase activity corresponds to amino acid sequences selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9,  
15 SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12 and wherein the amino acid sequences encompasses amino acid substitutions, deletions or insertions that do not alter the functional properties of the enzyme.

13. The method of Claim 1 wherein the protein having glycerol-3-phosphatase activity corresponds to the amino acid sequences selected from the group consisting of SEQ ID NO:13 and SEQ ID NO:14, and wherein the amino acid sequences may encompass amino acid substitutions, deletions or additions that do not alter the function of the enzyme.

14. A transformed host cell comprising:

- (a) a gene encoding a protein having a glycerol-3-phosphate  
25 dehydrogenase activity;
- (b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity;
- (c) a disruption in a gene encoding an endogenous glycerol kinase and;
- (d) a disruption a gene encoding an endogenous glycerol  
30 dehydrogenase;

wherein the disruptions in the genes of (c) and (d) prevent the expression of active gene product, and wherein the host cell converts at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides,  
35 polysaccharides, and single-carbon substrates to glycerol.

15. A transformed host cell comprising:

- (a) a gene encoding a protein having a glycerol-3-phosphate dehydrogenase activity;

(b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity; and

(c) a disruption in a gene encoding an endogenous glycerol dehydrogenase;

5 wherein the disruption in the gene of (c) prevents the expression of active gene product, and wherein the host cell converts at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and single-carbon substrates to glycerol.

16. A transformed host cell comprising:

10 (a) a gene encoding a protein having a glycerol-3-phosphate dehydrogenase activity;

(b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity; and

15 (c) a disruption in a gene encoding an endogenous glycerol kinase,

wherein the disruption in the gene of (c) prevents the expression of active gene product, and wherein the host cell converts at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and single-carbon substrates to glycerol.

20 17. A method for the production of 1,3-propanediol from a recombinant organism comprising:

(i) transforming a suitable host cell with an expression cassette comprising either one or both of

25 (a) a gene encoding a protein having glycerol-3-phosphate dehydrogenase activity, and

(b) a gene encoding a protein having glycerol-3-phosphate phosphatase activity,

the suitable host cell having at least one gene encoding a protein having a dehydratase activity and having a disruption in either one or both of:

30 (a) an endogenous gene encoding a polypeptide having glycerol kinase activity, and

(b) an endogenous gene encoding a polypeptide having glycerol dehydrogenase activity,

35 wherein the disruption in the genes of (a) or (b) prevents the expression of active gene product;

(ii) culturing the transformed host cell of (i) in the presence of at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, and single-carbon substrates whereby 1,3-propanediol is produced; and

(iii) recovering the 1,3-propanediol produced in (ii).

18. The method of Claim 17 wherein the protein having a dehydratase activity is selected from the group consisting of a glycerol dehydratase enzyme and a diol dehydratase enzyme.

5        19. The method of Claim 18 wherein the glycerol dehydratase enzyme is encoded by a gene, the gene isolated from a microorganism, the microorganism selected from the group consisting of *Klebsiella*, *Lactobacillus*, *Enterobacter*, *Citrobacter*, *Pelobacter*, *Ilyobacter*, and *Clostridium*.

10       20. The method of Claim 18 wherein the diol dehydratase enzyme is encoded by a gene, the gene isolated from a microorganism, the microorganism selected from the group consisting of *Klebsiella* and *Salmonella*.



SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT:
  - (A) NAME: E. I. DU PONT DE NEMOURS AND COMPANY
  - (B) STREET: 1007 MARKET STREET
  - (C) CITY: WILMINGTON
  - (D) STATE: DELAWARE
  - (E) COUNTRY: U.S.A.
  - (F) ZIP: 19898
  - (G) TELEPHONE: 302-892-8112
  - (H) TELEFAX: 302-773-0164
  - (I) TELEX: 6717325
- (ii) TITLE OF INVENTION: METHOD FOR THE PRODUCTION OF GLYCEROL BY RECOMBINANT ORGANISMS
- (iii) NUMBER OF SEQUENCES: 43
- (iv) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: DISKETTE, 3.5 INCH
  - (B) COMPUTER: IBM PC COMPATIBLE
  - (C) OPERATING SYSTEM: MICROSOFT WINDOWS 95
  - (D) SOFTWARE: MICROSOFT WORD VERSION 7.0A
- (v) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vi) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 08/982,783
  - (B) FILING DATE: DECEMBER 2, 1997
  - (C) CLASSIFICATION:
- (vii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: FLOYD, LINDA AXAMETHY
  - (B) REGISTRATION NUMBER: 33,692
  - (C) REFERENCE/DOCKET NUMBER: CR-9981-C

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1380 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```
CTTTAATTTT CTTTATCTT ACTCTCCTAC ATAAGACATC AAGAAACAAT TGTATATTGT    60
ACACCCCCCC CCTCCACAAA CACAAATATT GATAATATAA AGATGTCTGC TGCTGCTGAT    120
AGATTAAACT TAACTTCCGG CCACTTGAAT GCTGGTAGAA AGAGAAGTTC CTCTTCTGTT    180
TCTTTGAAGG CTGCCGAAAA GCCTTTCAAG GTTACTGTGA TTGGATCTGG TAACTGGGGT    240
ACTACTATTG CCAAGGTGGT TGCCGAAAAT TGTAAGGGAT ACCCAGAAGT TTTCGCTCCA    300
ATAGTACAAA TGTGGGTGTT CGAAGAAGAG ATCAATGGTG AAAAATTGAC TGAAATCATA    360
AATACTAGAC ATCAAAACGT GAAATACTTG CCTGGCATCA CTCTACCCGA CAATTTGGTT    420
GCTAATCCAG ACTTGATTGA TTCAGTCAAG GATGTCGACA TCATCGTTTT CAACATTCCA    480
CATCAATTTT TGCCCCGTAT CTGTAGCCAA TTGAAAGGTC ATGTTGATTC ACACGTCAGA    540
GCTATCTCCT GTCTAAAGGG TTTTGAAGTT GGTGCTAAAG GTGTCCAATT GCTATCCTCT    600
TACATCACTG AGGAACTAGG TATTCAATGT GGTGCTCTAT CTGGTGCTAA CATTGCCACC    660
GAAGTCGCTC AAGAACACTG GTCTGAAACA ACAGTTGCTT ACCACATTCC AAAGGATTTC    720
AGAGGCGAGG GCAAGGACGT CGACCATAAG GTTCTAAAGG CCTTGTTCCA CAGACCTTAC    780
TTCCACGTTA GTGTCATCGA AGATGTTGCT GGTATCTCCA TCTGTGGTGC TTTGAAGAAC    840
GTTGTTGCCT TAGGTTGTGG TTTCGTCGAA GGTCTAGGCT GGGGTAACAA CGCTTCTGCT    900
GCCATCCAAA GAGTCGGTTT GGGTGAGATC ATCAGATTCTG GTCAAATGTT TTTCCCAGAA    960
TCTAGAGAAG AAACATACTA CCAAGAGTCT GCTGGTGTTG CTGATTTGAT CACCACCTGC   1020
GCTGGTGGTA GAAACGTCAA GGTTGCTAGG CTAATGGCTA CTTCTGGTAA GGACGCCTGG   1080
GAATGTGAAA AGGAGTTGTT GAATGGCCAA TCCGCTCAAG GTTTAATTAC CTGCAAAGAA   1140
GTTACGAAT  GGTTGGAAAC ATGTGGCTCT GTCGAAGACT TCCCATTATT TGAAGCCGTA   1200
TACCAAATCG TTTACAACAA CTACCCAATG AAGAACCTGC CGGACATGAT TGAAGAATTA   1260
GATCTACATG AAGATTAGAT TTATTGGAGA AAGATAACAT ATCATACTTC CCCCCTTTT   1320
TTCGAGGCTC TTCTATATCA TATTCATAAA TTAGCATTAT GTCATTTCTC ATAACACTTT   1380
```

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 2946 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```
GAATTCGAGC CTGAAGTGCT GATTACCTTC AGGTAGACTT CATCTTGACC CATCAACCCC      60
AGCGTCAATC CTGCAAATAC ACCACCCAGC AGCACTAGGA TGATAGAGAT AATATAGTAC      120
GTGGTAACGC TTGCCTCATC ACCTACGCTA TGGCCGGAAT CGGCAACATC CCTAGAATTG      180
AGTACGTGTG ATCCGGATAA CAACGGCAGT GAATATATCT TCGGTATCGT AAAGATGTGA      240
TATAAGATGA TGTATACCCA ATGAGGAGCG CCTGATCGTG ACCTAGACCT TAGTGGCAAA      300
AACGACATAT CTATTATAGT GGGGAGAGTT TCGTGCAAAT AACAGACGCA GCAGCAAGTA      360
ACTGTGACGA TATCAACTCT TTTTTTATTA TGTAATAAGC AAACAAGCAC GAATGGGGAA      420
AGCCTATGTG CAATCACCAA GGTCGTCCCT TTTTCCCAT TTGCTAATTT AGAATTTAAA      480
GAAACCAAAA GAATGAAGAA AGAAAACAAA TACTAGCCCT AACCTGACT TCGTTTCTAT      540
GATAATACCC TGCTTTAATG AACGGTATGC CCTAGGGTAT ATCTCACTCT GTACGTTACA      600
AACTCCGGTT ATTTTATCGG AACATCCGAG CACCCGCGCC TTCCTCAACC CAGGCACCGC      660
CCCAGGTAAC CGTGCGCGAT GAGCTAATCC TGAGCCATCA CCCACCCAC CCGTTGATGA      720
CAGCAATTCTG GGAGGGCGAA AATAAACTG GAGCAAGGAA TTACCATCAC CGTCACCATC      780
ACCATCATAT CGCCTTAGCC TCTAGCCATA GCCATCATGC AAGCGTGTAT CTTCTAAGAT      840
TCAGTCATCA TCATTACCGA GTTTGTTTTT CTCACATGA TGAAGAAGGT TTGAGTATGC      900
TCGAAACAAT AAGACGACGA TGGCTCTGCC ATTGGTTATA TTACGCTTTT GCGGCGAGGT      960
GCCGATGGGT TGCTGAGGGG AAGAGTGTTT AGCTTACGGA CCTATTGCCA TTGTTATTCC     1020
GATTAATCTA TTGTTTACGA GCTCTTCTCT ACCCTGTCAT TCTAGTATTT TTTTTTTTTT     1080
TTTTTGTTTT TACTTTTTTT TCTTCTTGCC TTTTTTCTT GTTACTTTTT TTCTAGTTTT     1140
TTTTCCTTCC ACTAAGCTTT TTCCTTGATT TATCCTTGGG TTCTTCTTTC TACTCCTTTA     1200
GATTTTTTTT TTATATATTA ATTTTAAAGT TTATGTATTT TGGTAGATTC AATTCTCTTT     1260
CCCTTTCCTT TTCCTTCGCT CCCCTTCCTT ATCAATGCTT GCTGTCAGAA GATTAACAAG     1320
ATACACATTC CTTAAGCGAA CGCATCCGGT GTTATATACT CGTCGTGCAT ATAAAATTTT     1380
```

GCCTTCAAGA TCTACTTTCC TAAGAAGATC ATTATTACAA ACACAACTGC ACTCAAAGAT	1440
GACTGCTCAT ACTAATATCA AACAGCACAA AACTGTGCAT GAGGACCATC CTATCAGAAG	1500
ATCGGACTCT GCCGTGTCAA TTGTACATTT GAAACGTGCG CCCTTCAAGG TTACAGTGAT	1560
TGGTTCTGGT AACTGGGGGA CCACCATCGC CAAAGTCATT GCGGAAAACA CAGAATTGCA	1620
TTCCCATATC TTCGAGCCAG AGGTGAGAAT GTGGGTTTTT GATGAAAAGA TCGGCGACGA	1680
AAATCTGACG GATATCATAA ATACAAGACA CCAGAACGTT AAATATCTAC CCAATATTGA	1740
CCTGCCCCAT AATCTAGTGG CCGATCCTGA TCTTTTACAC TCCATCAAGG GTGCTGACAT	1800
CCTTGTTTTT AACATCCCTC ATCAATTTTT ACCAAACATA GTCAAACAAT TGCAAGGCCA	1860
CGTGGCCCCCT CATGTAAGGG CCATCTCGTG TCTAAAAGGG TTCGAGTTGG GCTCCAAGGG	1920
TGTGCAATTG CTATCCTCCT ATGTTACTGA TGAGTTAGGA ATCCAATGTG GCGCACTATC	1980
TGGTGCAAAC TTGGCACCGG AAGTGGCCAA GGAGCATTGG TCCGAAACCA CCGTGGCTTA	2040
CCAAC TACCA AAGGATTATC AAGGTGATGG CAAGGATGTA GATCATAAGA TTTTGAAATT	2100
GCTGTTCCAC AGACCTTACT TCCACGTCAA TGTCATCGAT GATGTTGCTG GTATATCCAT	2160
TGCCGGTGCC TTGAAGAACG TCGTGGCACT TGCATGTGGT TTCGTAGAAG GTATGGGATG	2220
GGGTAACAAT GCCTCCGCAG CCATTCAAAG GCTGGGTTTA GGTGAAATTA TCAAGTTCGG	2280
TAGAATGTTT TTCCCAGAAT CCAAAGTCGA GACCTACTAT CAAGAATCCG CTGGTGTTGC	2340
AGATCTGATC ACCACCTGCT CAGGCGGTAG AAACGTCAAG GTTGCCACAT ACATGGCCAA	2400
GACCGGTAAG TCAGCCTTGG AAGCAGAAAA GGAATTGCTT AACGGTCAAT CCGCCCAAGG	2460
GATAATCACA TGCAGAGAAG TTCACGAGTG GCTACAAACA TGTGAGTTGA CCCAAGAATT	2520
CCCAATTATT CGAGGCAGTC TACCAGATAG TCTACAACAA CGTCCGCATG GAAGACCTAC	2580
CGGAGATGAT TGAAGAGCTA GACATCGATG ACGAATAGAC ACTCTCCCCC CCCCTCCCCC	2640
TCTGATCTTT CCTGTTGCCT CTTTTTCCCC CAACCAATTT ATCATTATAC ACAAGTTCTA	2700
CAACTACTAC TAGTAACATT ACTACAGTTA TTATAATTTT CTATTCTCTT TTTCTTTAAG	2760
AATCTATCAT TAACGTTAAT TTCTATATAT ACATAACTAC CATTATACAC GCTATTATCG	2820
TTTACATATC ACATCACCGT TAATGAAAGA TACGACACCC TGTACACTAA CACAATTAAA	2880
TAATCGCCAT AACCTTTTCT GTTATCTATA GCCCTTAAAG CTGTTTCTTC GAGCTTTTCA	2940
CTGCAG	2946

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3178 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

CTGCAGAACT TCGTCTGCTC TGTGCCCATC CTCGCGGTTA GAAAGAAGCT GAATTGTTTC    60
ATGCGCAAGG GCATCAGCGA GTGACCAATA ATCACTGCAC TAATTCCTTT TTAGCAACAC    120
ATACTTATAT ACAGCACCAG ACCTTATGTC TTTTCTCTGC TCCGATACGT TATCCCACCC    180
AACTTTTATT TCAGTTTTTG CAGGGGAAAT TTCACAACCC CGCACGCTAA AAATCGTATT    240
TAAACTTAAA AGAGAACAGC CACAAATAGG GAACTTTGGT CTAAACGAAG GACTCTCCCT    300
CCCTTATCTT GACCGTGCTA TTGCCATCAC TGCTACAAGA CTAAATACGT ACTAATATAT    360
GTTTTCGGTA ACGAGAAGAA GAGCTGCCGG TGCAGCTGCT GCCATGGCCA CAGCCACGGG    420
GACGCTGTAC TGGATGACTA GCCAAGGTGA TAGGCCGTTA GTGCACAATG ACCCGAGCTA    480
CATGGTGCAA TTCCCCACCG CCGCTCCACC GGCAGGTCTC TAGACGAGAC CTGCTGGACC    540
GTCTGGACAA GACGCATCAA TTCGACGTGT TGATCATCGG TGGCGGGGCC ACGGGGACAG    600
GATGTGCCCT AGATGCTGCG ACCAGGGGAC TCAATGTGGC CTTTGTGAA AAGGGGGATT    660
TTGCCTCGGG AACGTCGTCC AAATCTACCA AGATGATTCA CGGTGGGGTG CGGTACTTAG    720
AGAAGGCCTT CTGGGAGTTC TCCAAGGCAC AACTGGATCT GGTCATCGAG GCACTCAACG    780
AGCGTAAACA TCTTATCAAC ACTGCCCCTC ACCTGTGCAC GGTGCTACCA ATTCTGATCC    840
CCATCTACAG CACCTGGCAG GTCCCGTACA TCTATATGGG CTGTAAATTC TACGATTTCT    900
TTGGCGGTTC CCAAACCTTG AAAAAATCAT ACCTACTGTC CAAATCCGCC ACCGTGGAGA    960
AGGCTCCCAT GCTTACCACA GACAATTTAA AGGCCTCGCT TGTGTACCAT GATGGGTCCT   1020
TTAACGACTC GCGTTTGAAC GCCACTTTAG CCATCACGGG TGTGGAGAAC GGCGCTACCG   1080
TCTTGATCTA TGTCGAGGTA CAAAATTGA TCAAAGACCC AACTTCTGGT AAGGTTATCG   1140
GTGCCGAGGC CCGGGACGTT GAGACTAATG AGCTTGTCAG AATCAACGCT AAATGTGTGG   1200
TCAATGCCAC GGGCCCATAC AGTGACGCCA TTTTGCAAAT GGACCGCAAC CCATCCGGTC   1260
TGCCGGACTC CCCGCTAAAC GACAACTCCA AGATCAAGTC GACTTTCAAT CAAATCTCCG   1320
TCATGGACCC GAAAATGGTC ATCCCATCTA TTGGCGTTCA CATCGTATTG CCCTCTTTTT   1380

```



ACTCCCCGAA GGATATGGGT TTGTTGGACG TCAGAACCTC TGATGGCAGA GTGATGTTCT 1440  
TTTTACCTTG GCAGGGCAAA GTCCTTGCCG GCACCACAGA CATCCCCTA AAGCAAGTCC 1500  
CAGAAAACCC TATGCCTACA GAGGCTGATA TTCAAGATAT CTTGAAAGAA CTACAGCACT 1560  
ATATCGAATT CCCCCTGAAA AGAGAAGACG TGCTAAGTGC ATGGGCTGGT GTCAGACCTT 1620  
TGGTCAGAGA TCCACGTACA ATCCCCGCAG ACGGGAAGAA GGGCTCTGCC ACTCAGGGCG 1680  
TGGTAAGATC CCACTTCTTG TTCACTTCGG ATAATGGCCT AATTACTATT GCAGGTGGTA 1740  
AATGGACTAC TTACAGACAA ATGGCTGAGG AAACAGTCGA CAAAGTTGTC GAAGTTGGCG 1800  
GATTCCACAA CCTGAAACCT TGTCACACAA GAGATATTAA GCTTGCTGGT GCAGAAGAAT 1860  
GGACGCAAAA CTATGTGGCT TTATTGGCTC AAAACTACCA TTTATCATCA AAAATGTCCA 1920  
ACTACTTGGT TCAAACTAC GGAACCCGTT CCTCTATCAT TTGCGAATTT TTCAAAGAAT 1980  
CCATGGAAAA TAAACTGCCT TTGTCCTTAG CCGACAAGGA AAATAACGTA ATCTACTCTA 2040  
GCGAGGAGAA CAACTTGGTC AATTTTGATA CTTTCAGATA TCCATTCACA ATCGGTGAGT 2100  
TAAAGTATTC CATGCAGTAC GAATATTGTA GAACTCCCTT GGAATTCCTT TTAAGAAGAA 2160  
CAAGATTCGC CTTCTTGGAC GCCAAGGAAG CTTTGAATGC CGTGCATGCC ACCGTCAAAG 2220  
TTATGGGTGA TGAGTTCAAT TGGTCGGAGA AAAAGAGGCA GTGGGAACTT GAAAAAACTG 2280  
TGAACTTCAT CCAAGGACGT TTCGGTGTCT AAATCGATCA TGATAGTTAA GGGTGACAAA 2340  
GATAACATTC ACAAGAGTAA TAATAATGGT AATGATGATA ATAATAATAA TGATAGTAAT 2400  
AACAATAATA ATAATGGTGG TAATGGCAAT GAAATCGCTA TTATTACCTA TTTTCCTTAA 2460  
TGGAAGAGTT AAAGTAACT AAAAAAACTA CAAAAATATA TGAAGAAAAA AAAAAAAGA 2520  
GGTAATAGAC TCTACTACTA CAATTGATCT TCAAATTATG ACCTTCCTAG TGTTTATATT 2580  
CTATTTCCAA TACATAATAT AATCTATATA ATCATTGCTG GTAGACTTCC GTTTTAATAT 2640  
CGTTTTAATT ATCCCCTTTA TCTCTAGTCT AGTTTTATCA TAAATATAG AAACACTAAA 2700  
TAATATTCTT CAAACGGTCC TGGTGCATAC GCAATACATA TTTATGGTGC AAAAAAAAAA 2760  
ATGGAAAATT TTGCTAGTCA TAAACCCTTT CATAAAACAA TACGTAGACA TCGCTACTTG 2820  
AAATTTTCAA GTTTTTATCA GATCCATGTT TCCTATCTGC CTTGACAACC TCATCGTCGA 2880  
AATAGTACCA TTTAGAACGC CCAATATTCA CATTGTGTTT AAGGTCTTTA TTCACCAGTG 2940  
ACGTGTAATG GCCATGATTA ATGTGCCTGT ATGGTTAACC ACTCCAAATA GCTTATATTT 3000  
CATAGTGTCA TTGTTTTTCA ATATAATGTT TAGTATCAAT GGATATGTTA CGACGGTGTT 3060  
ATTTTTCTTG GTCAAATCGT AATAAAATCT CGATAAATGG ATGACTAAGA TTTTGGTAA 3120

AGTTACAAAA TTTATCGTTT TCACTGTTGT CAATTTTTTG TTCTTGTAAT CACTCGAG 3178

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 816 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATGAAACGTT TCAATGTTTT AAAATATATC AGAACAACAA AAGCAAATAT ACAAACCATC	60
GCAATGCCTT TGACCACAAA ACCTTTATCT TTGAAAATCA ACGCCGCTCT ATTCGATGTT	120
GACGGTACCA TCATCATCTC TCAACCAGCC ATTGCTGCTT TCTGGAGAGA TTTCGGTAAA	180
GACAAGCCTT ACTTCGATGC CGAACACGTT ATTCACATCT CTCACGGTTG GAGAACTTAC	240
GATGCCATTG CCAAGTTCGC TCCAGACTTT GCTGATGAAG AATACGTTAA CAAGCTAGAA	300
GGTGAAATCC CAGAAAAGTA CGGTGAACAC TCCATCGAAG TTCCAGGTGC TGTCAGTTG	360
TGTAATGCTT TGAACGCCTT GCCAAAGGAA AAATGGGCTG TCGCCACCTC TGGTACCCGT	420
GACATGGCCA AGAAATGGTT CGACATTTTG AAGATCAAGA GACCAGAATA CTTCATCACC	480
GCCAATGATG TCAAGCAAGG TAAGCCTCAC CCAGAACCAT ACTTAAAGGG TAGAAACGGT	540
TTGGGTTTCC CAATTAATGA ACAAGACCCA TCCAAATCTA AGGTTGTTGT CTTTGAAGAC	600
GCACCAGCTG GTATTGCTGC TGGTAAGGCT GCTGGCTGTA AAATCGTTGG TATTGCTACC	660
ACTTTCGATT TGGACTTCTT GAAGGAAAAG GGTGTGACA TCATTGTCAA GAACCACGAA	720
TCTATCAGAG TCGGTGAATA CAACGCTGAA ACCGATGAAG TCGAATTGAT CTTTGATGAC	780
TACTTATACG CTAAGGATGA CTTGTTGAAA TGGTAA	816

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 753 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATGGGATTGA CTACTAAACC TCTATCTTTG AAAGTTAACG CCGCTTTGTT CGACGTCGAC	60
GGTACCATTA TCATCTCTCA ACCAGCCATT GCTGCATTCT GGAGGGATTT CGGTAAGGAC	120

AAACCTTATT TCGATGCTGA ACACGTTATC CAAGTCTCGC ATGGTTGGAG AACGTTTGAT 180  
 GCCATTGCTA AGTTCGCTCC AGACTTTGCC AATGAAGAGT ATGTTAACAA ATTAGAAGCT 240  
 GAAATTCCGG TCAAGTACGG TGAAAAATCC ATTGAAGTCC CAGGTGCAGT TAAGCTGTGC 300  
 AACGCTTTGA ACGCTCTACC AAAAGAGAAA TGGGCTGTGG CAACTTCCGG TACCCGTGAT 360  
 ATGGCACAAA AATGGTTCGA GCATCTGGGA ATCAGGAGAC CAAAGTACTT CATTACCGCT 420  
 AATGATGTCA AACAGGGTAA GCCTCATCCA GAACCATATC TGAAGGGCAG GAATGGCTTA 480  
 GGATATCCGA TCAATGAGCA AGACCCTTCC AAATCTAAGG TAGTAGTATT TGAAGACGCT 540  
 CCAGCAGGTA TTGCCGCCGG AAAAGCCGCC GGTGTGAAGA TCATTGGTAT TGCCACTACT 600  
 TTCGACTTGG ACTTCCTAAA GGAAAAAGGC TGTGACATCA TTGTCAAAAA CCACGAATCC 660  
 ATCAGAGTTG GCGGCTACAA TGCCGAAACA GACGAAGTTG AATTCATTTT TGACGACTAC 720  
 TTATATGCTA AGGACGATCT GTTGAAATGG TAA 753

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2520 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TGTATTGGCC ACGATAACCA CCCTTTGTAT ACTGTTTTTG TTTTTCACAT GGTAATAAAC 60  
 GACTTTTATT AAACAACGTA TGTA AAAACA TAACAAGAAT CTACCCATAC AGGCCATTTC 120  
 GTAATTCTTC TCTTCTAATT GGAGTAAAAC CATCAATTAA AGGGTGTGGA GTAGCATAGT 180  
 GAGGGGCTGA CTGCATTGAC AAAAAAATTG AAAAAAAAAA AGGAAAAGGA AAGGAAAAAA 240  
 AGACAGCCAA GACTTTTAGA ACGGATAAGG TGTAATAAAA TGTGGGGGGA TGCCTGTTCT 300  
 CGAACCATAT AAAATATAACC ATGTGGTTTG AGTTGTGGCC GGAACATAC AAATAGTTAT 360  
 ATGTTTCCCT CTCTCTTCCG ACTTGTAGTA TTCTCCAAAC GTTACATATT CCGATCAAGC 420  
 CAGCGCCTTT AACTAGTTT AAAACAAGAA CAGAGCCGTA TGTCCAAAAT AATGGAAGAT 480  
 TTACGAAGTG ACTACGTCCC GCTTATCGCC AGTATTGATG TAGGAACGAC CTCATCCAGA 540  
 TGCATTCTGT TCAACAGATG GGGCCAGGAC GTTTCAAAAC ACCAAATTGA ATATTCAACT 600  
 TCAGCATCGA AGGGCAAGAT TGGGGTGTCT GGCCTAAGGA GACCCTCTAC AGCCCCAGCT 660  
 CGTGAAACAC CAAACGCCGG TGACATCAAA ACCAGCGGAA AGCCCATCTT TTCTGCAGAA 720

GGCTATGCCA TTCAAGAAAC CAAATTCCTA AAAATCGAGG AATTGGACTT GGACTTCCAT 780  
AACGAACCCA CGTTGAAGTT CCCCAAACCG GGTGGGTTG AGTGCCATCC GCAGAAATTA 840  
CTGGTGAACG TCGTCCAATG CCTTGCCTCA AGTTTGCTCT CTCTGCAGAC TATCAACAGC 900  
GAACGTGTAG CAAACGGTCT CCCACCTTAC AAGGTAATAT GCATGGGTAT AGCAAACATG 960  
AGAGAAACCA CAATTCTGTG GTCCCGCCGC ACAGGAAAAC CAATTGTTAA CTACGGTATT 1020  
GTTTGGAACG ACACCAGAAC GATCAAAATC GTTAGAGACA AATGGCAAAA CACTAGCGTC 1080  
GATAGGCAAC TGCAGCTTAG ACAGAAGACT GGATTGCCAT TGCTCTCCAC GTATTTCTCC 1140  
TGTTCCAAGC TCGCTGGTT CCTCGACAAT GAGCCTCTGT GTACCAAGGC GTATGAGGAG 1200  
AACGACCTGA TGTTGGGCAC TGTGGACACA TGGCTGATTT ACCAATTAAC TAAACAAAAG 1260  
GCGTTCGTTT CTGACGTAAC CAACGCTTCC AGAACTGGAT TTATGAACCT CTCCACTTTA 1320  
AAGTACGACA ACGAGTTGCT GGAATTTTGG GGTATTGACA AGAACCTGAT TCACATGCCC 1380  
GAAATTGTGT CCTCATCTCA ATACTACGGT GACTTTGGCA TTCCTGATTG GATAATGGAA 1440  
AAGCTACACG ATTCGCCAAA AACAGTACTG CGAGATCTAG TCAAGAGAAA CCTGCCATA 1500  
CAGGGCTGTC TGGGCGACCA AAGCGCATCC ATGGTGGGGC AACTCGCTTA CAAACCCGGT 1560  
GCTGCAAAAT GTACTTATGG TACCGGTTGC TTTTACTGT ACAATACGGG GACCAAAAAA 1620  
TTGATCTCCC AACATGGCGC ACTGACGACT CTAGCATTTT GGTTCACACA TTTGCAAGAG 1680  
TACGGTGGCC AAAAACCAGA ATTGAGCAAG CCACATTTTG CATTAGAGGG TTCCGTCGCT 1740  
GTGGCTGGTG CTGTGGTCCA ATGGCTACGT GATAATTTAC GATTGATCGA TAAATCAGAG 1800  
GATGTCGGAC CGATTGCATC TACGGTTCCT GATTCTGGTG GCGTAGTTTT CGTCCCCGCA 1860  
TTTAGTGGCC TATTCGCTCC CTATTGGGAC CCAGATGCCA GAGCCACCAT AATGGGGATG 1920  
TCTCAATTCA CTACTGCCTC CCACATCGCC AGAGCTGCCG TGGAAGGTGT TTGCTTTCAA 1980  
GCCAGGGCTA TCTTGAAGGC AATGAGTTCT GACGCGTTTG GTGAAGGTTC CAAAGACAGG 2040  
GACTTTTTAG AGGAAATTTT CGACGTCACA TATGAAAAGT CGCCCCTGTC GGTCTGGCA 2100  
GTGGATGGCG GGATGTCGAG GTCTAATGAA GTCATGCAAA TTCAAGCCGA TATCCTAGGT 2160  
CCCTGTGTCA AAGTCAGAAG GTCTCCGACA GCGGAATGTA CCGCATTGGG GGCAGCCATT 2220  
GCAGCCAATA TGGCTTTCAA GGATGTGAAC GAGCGCCCAT TATGGAAGGA CCTACACGAT 2280  
GTTAAGAAAT GGGTCTTTTA CAATGGAATG GAGAAAAACG AACAAATATC ACCAGAGGCT 2340  
CATCCAAACC TTAAGATATT CAGAAGTGAA TCCGACGATG CTGAAAGGAG AAAGCATTGG 2400  
AAGTATTGGG AAGTTGCCGT GGAAAGATCC AAAGGTTGGC TGAAGGACAT AGAAGGTGAA 2460

CACGAACAGG TTCTAGAAAA CTTCCAATAA CAACATAAAT AATTTCTATT AACAATGTAA 2520

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 391 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Ser	Ala	Ala	Ala	Asp	Arg	Leu	Asn	Leu	Thr	Ser	Gly	His	Leu	Asn	1	5	10	15
Ala	Gly	Arg	Lys	Arg	Ser	Ser	Ser	Ser	Val	Ser	Leu	Lys	Ala	Ala	Glu	20	25	30	
Lys	Pro	Phe	Lys	Val	Thr	Val	Ile	Gly	Ser	Gly	Asn	Trp	Gly	Thr	Thr	35	40	45	
Ile	Ala	Lys	Val	Val	Ala	Glu	Asn	Cys	Lys	Gly	Tyr	Pro	Glu	Val	Phe	50	55	60	
Ala	Pro	Ile	Val	Gln	Met	Trp	Val	Phe	Glu	Glu	Glu	Ile	Asn	Gly	Glu	65	70	75	80
Lys	Leu	Thr	Glu	Ile	Ile	Asn	Thr	Arg	His	Gln	Asn	Val	Lys	Tyr	Leu	85	90	95	
Pro	Gly	Ile	Thr	Leu	Pro	Asp	Asn	Leu	Val	Ala	Asn	Pro	Asp	Leu	Ile	100	105	110	
Asp	Ser	Val	Lys	Asp	Val	Asp	Ile	Ile	Val	Phe	Asn	Ile	Pro	His	Gln	115	120	125	
Phe	Leu	Pro	Arg	Ile	Cys	Ser	Gln	Leu	Lys	Gly	His	Val	Asp	Ser	His	130	135	140	
Val	Arg	Ala	Ile	Ser	Cys	Leu	Lys	Gly	Phe	Glu	Val	Gly	Ala	Lys	Gly	145	150	155	160
Val	Gln	Leu	Leu	Ser	Ser	Tyr	Ile	Thr	Glu	Glu	Leu	Gly	Ile	Gln	Cys	165	170	175	
Gly	Ala	Leu	Ser	Gly	Ala	Asn	Ile	Ala	Thr	Glu	Val	Ala	Gln	Glu	His	180	185	190	
Trp	Ser	Glu	Thr	Thr	Val	Ala	Tyr	His	Ile	Pro	Lys	Asp	Phe	Arg	Gly	195	200	205	
Glu	Gly	Lys	Asp	Val	Asp	His	Lys	Val	Leu	Lys	Ala	Leu	Phe	His	Arg	210	215	220	
Pro	Tyr	Phe	His	Val	Ser	Val	Ile	Glu	Asp	Val	Ala	Gly	Ile	Ser	Ile	225	230	235	240



(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Thr	Ala	His	Thr	Asn	Ile	Lys	Gln	His	Lys	His	Cys	His	Glu	Asp
1				5					10					15	
His	Pro	Ile	Arg	Arg	Ser	Asp	Ser	Ala	Val	Ser	Ile	Val	His	Leu	Lys
			20					25					30		
Arg	Ala	Pro	Phe	Lys	Val	Thr	Val	Ile	Gly	Ser	Gly	Asn	Trp	Gly	Thr
		35					40					45			
Thr	Ile	Ala	Lys	Val	Ile	Ala	Glu	Asn	Thr	Glu	Leu	His	Ser	His	Ile
	50					55					60				
Phe	Glu	Pro	Glu	Val	Arg	Met	Trp	Val	Phe	Asp	Glu	Lys	Ile	Gly	Asp
65					70					75					80

Glu Asn Leu Thr Asp Ile Ile Asn Thr Arg His Gln Asn Val Lys Tyr  
 85 90 95  
 Leu Pro Asn Ile Asp Leu Pro His Asn Leu Val Ala Asp Pro Asp Leu  
 100 105 110  
 Leu His Ser Ile Lys Gly Ala Asp Ile Leu Val Phe Asn Ile Pro His  
 115 120 125  
 Gln Phe Leu Pro Asn Ile Val Lys Gln Leu Gln Gly His Val Ala Pro  
 130 135 140  
 His Val Arg Ala Ile Ser Cys Leu Lys Gly Phe Glu Leu Gly Ser Lys  
 145 150 155 160  
 Gly Val Gln Leu Leu Ser Ser Tyr Val Thr Asp Glu Leu Gly Ile Gln  
 165 170 175  
 Cys Gly Ala Leu Ser Gly Ala Asn Leu Ala Pro Glu Val Ala Lys Glu  
 180 185 190  
 His Trp Ser Glu Thr Thr Val Ala Tyr Gln Leu Pro Lys Asp Tyr Gln  
 195 200 205  
 Gly Asp Gly Lys Asp Val Asp His Lys Ile Leu Lys Leu Leu Phe His  
 210 215 220  
 Arg Pro Tyr Phe His Val Asn Val Ile Asp Asp Val Ala Gly Ile Ser  
 225 230 235 240  
 Ile Ala Gly Ala Leu Lys Asn Val Val Ala Leu Ala Cys Gly Phe Val  
 245 250 255  
 Glu Gly Met Gly Trp Gly Asn Asn Ala Ser Ala Ala Ile Gln Arg Leu  
 260 265 270  
 Gly Leu Gly Glu Ile Ile Lys Phe Gly Arg Met Phe Phe Pro Glu Ser  
 275 280 285  
 Lys Val Glu Thr Tyr Tyr Gln Glu Ser Ala Gly Val Ala Asp Leu Ile  
 290 295 300  
 Thr Thr Cys Ser Gly Gly Arg Asn Val Lys Val Ala Thr Tyr Met Ala  
 305 310 315 320  
 Lys Thr Gly Lys Ser Ala Leu Glu Ala Glu Lys Glu Leu Leu Asn Gly  
 325 330 335  
 Gln Ser Ala Gln Gly Ile Ile Thr Cys Arg Glu Val His Glu Trp Leu  
 340 345 350  
 Gln Thr Cys Glu Leu Thr Gln Glu Phe Pro Ile Ile Arg Gly Ser Leu  
 355 360 365  
 Pro Asp Ser Leu Gln Gln Arg Pro His Gly Arg Pro Thr Gly Asp Asp  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 614 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Met Thr Arg Ala Thr Trp Cys Asn Ser Pro Pro Pro Leu His Arg Gln
1          5          10          15

Val Ser Arg Arg Asp Leu Leu Asp Arg Leu Asp Lys Thr His Gln Phe
          20          25          30

Asp Val Leu Ile Ile Gly Gly Gly Ala Thr Gly Thr Gly Cys Ala Leu
          35          40          45

Asp Ala Ala Thr Arg Gly Leu Asn Val Ala Leu Val Glu Lys Gly Asp
          50          55          60

Phe Ala Ser Gly Thr Ser Ser Lys Ser Thr Lys Met Ile His Gly Gly
65          70          75          80

Val Arg Tyr Leu Glu Lys Ala Phe Trp Glu Phe Ser Lys Ala Gln Leu
          85          90          95

Asp Leu Val Ile Glu Ala Leu Asn Glu Arg Lys His Leu Ile Asn Thr
          100          105          110

Ala Pro His Leu Cys Thr Val Leu Pro Ile Leu Ile Pro Ile Tyr Ser
          115          120          125

Thr Trp Gln Val Pro Tyr Ile Tyr Met Gly Cys Lys Phe Tyr Asp Phe
          130          135          140

Phe Gly Gly Ser Gln Asn Leu Lys Lys Ser Tyr Leu Leu Ser Lys Ser
145          150          155          160

Ala Thr Val Glu Lys Ala Pro Met Leu Thr Thr Asp Asn Leu Lys Ala
          165          170          175

Ser Leu Val Tyr His Asp Gly Ser Phe Asn Asp Ser Arg Leu Asn Ala
          180          185          190

Thr Leu Ala Ile Thr Gly Val Glu Asn Gly Ala Thr Val Leu Ile Tyr
          195          200          205

Val Glu Val Gln Lys Leu Ile Lys Asp Pro Thr Ser Gly Lys Val Ile
          210          215          220

Gly Ala Glu Ala Arg Asp Val Glu Thr Asn Glu Leu Val Arg Ile Asn
225          230          235          240

```

Ala	Lys	Cys	Val	Val	Asn	Ala	Thr	Gly	Pro	Tyr	Ser	Asp	Ala	Ile	Leu	245	250	255
Gln	Met	Asp	Arg	Asn	Pro	Ser	Gly	Leu	Pro	Asp	Ser	Pro	Leu	Asn	Asp	260	265	270
Asn	Ser	Lys	Ile	Lys	Ser	Thr	Phe	Asn	Gln	Ile	Ser	Val	Met	Asp	Pro	275	280	285
Lys	Met	Val	Ile	Pro	Ser	Ile	Gly	Val	His	Ile	Val	Leu	Pro	Ser	Phe	290	295	300
Tyr	Ser	Pro	Lys	Asp	Met	Gly	Leu	Leu	Asp	Val	Arg	Thr	Ser	Asp	Gly	305	310	315
Arg	Val	Met	Phe	Phe	Leu	Pro	Trp	Gln	Gly	Lys	Val	Leu	Ala	Gly	Thr	325	330	335
Thr	Asp	Ile	Pro	Leu	Lys	Gln	Val	Pro	Glu	Asn	Pro	Met	Pro	Thr	Glu	340	345	350
Ala	Asp	Ile	Gln	Asp	Ile	Leu	Lys	Glu	Leu	Gln	His	Tyr	Ile	Glu	Phe	355	360	365
Pro	Val	Lys	Arg	Glu	Asp	Val	Leu	Ser	Ala	Trp	Ala	Gly	Val	Arg	Pro	370	375	380
Leu	Val	Arg	Asp	Pro	Arg	Thr	Ile	Pro	Ala	Asp	Gly	Lys	Lys	Gly	Ser	385	390	395
Ala	Thr	Gln	Gly	Val	Val	Arg	Ser	His	Phe	Leu	Phe	Thr	Ser	Asp	Asn	405	410	415
Gly	Leu	Ile	Thr	Ile	Ala	Gly	Gly	Lys	Trp	Thr	Thr	Tyr	Arg	Gln	Met	420	425	430
Ala	Glu	Glu	Thr	Val	Asp	Lys	Val	Val	Glu	Val	Gly	Gly	Phe	His	Asn	435	440	445
Leu	Lys	Pro	Cys	His	Thr	Arg	Asp	Ile	Lys	Leu	Ala	Gly	Ala	Glu	Glu	450	455	460
Trp	Thr	Gln	Asn	Tyr	Val	Ala	Leu	Leu	Ala	Gln	Asn	Tyr	His	Leu	Ser	465	470	475
Ser	Lys	Met	Ser	Asn	Tyr	Leu	Val	Gln	Asn	Tyr	Gly	Thr	Arg	Ser	Ser	485	490	495
Ile	Ile	Cys	Glu	Phe	Phe	Lys	Glu	Ser	Met	Glu	Asn	Lys	Leu	Pro	Leu	500	505	510
Ser	Leu	Ala	Asp	Lys	Glu	Asn	Asn	Val	Ile	Tyr	Ser	Ser	Glu	Glu	Asn	515	520	525
Asn	Leu	Val	Asn	Phe	Asp	Thr	Phe	Arg	Tyr	Pro	Phe	Thr	Ile	Gly	Glu	530	535	540

Leu Lys Tyr Ser Met Gln Tyr Glu Tyr Cys Arg Thr Pro Leu Asp Phe  
545 550 555 560

Leu Leu Arg Arg Thr Arg Phe Ala Phe Leu Asp Ala Lys Glu Ala Leu  
565 570 575

Asn Ala Val His Ala Thr Val Lys Val Met Gly Asp Glu Phe Asn Trp  
580 585 590

Ser Glu Lys Lys Arg Gln Trp Glu Leu Glu Lys Thr Val Asn Phe Ile  
595 600 605

Gln Gly Arg Phe Gly Val  
610

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 339 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Asn Gln Arg Asn Ala Ser Met Thr Val Ile Gly Ala Gly Ser Tyr  
1 5 10 15

Gly Thr Ala Leu Ala Ile Thr Leu Ala Arg Asn Gly His Glu Val Val  
20 25 30

Leu Trp Gly His Asp Pro Glu His Ile Ala Thr Leu Glu Arg Asp Arg  
35 40 45

Cys Asn Ala Ala Phe Leu Pro Asp Val Pro Phe Pro Asp Thr Leu His  
50 55 60

Leu Glu Ser Asp Leu Ala Thr Ala Leu Ala Ala Ser Arg Asn Ile Leu  
65 70 75 80

Val Val Val Pro Ser His Val Phe Gly Glu Val Leu Arg Gln Ile Lys  
85 90 95

Pro Leu Met Arg Pro Asp Ala Arg Leu Val Trp Ala Thr Lys Gly Leu  
100 105 110

Glu Ala Glu Thr Gly Arg Leu Leu Gln Asp Val Ala Arg Glu Ala Leu  
115 120 125

Gly Asp Gln Ile Pro Leu Ala Val Ile Ser Gly Pro Thr Phe Ala Lys  
130 135 140

Glu Leu Ala Ala Gly Leu Pro Thr Ala Ile Ser Leu Ala Ser Thr Asp  
145 150 155 160



Gln Thr Phe Ala Asp Asp Leu Gln Gln Leu Leu His Cys Gly Lys Ser  
 165 170 175  
 Phe Arg Val Tyr Ser Asn Pro Asp Phe Ile Gly Val Gln Leu Gly Gly  
 180 185 190  
 Ala Val Lys Asn Val Ile Ala Ile Gly Ala Gly Met Ser Asp Gly Ile  
 195 200 205  
 Gly Phe Gly Ala Asn Ala Arg Thr Ala Leu Ile Thr Arg Gly Leu Ala  
 210 215 220  
 Glu Met Ser Arg Leu Gly Ala Ala Leu Gly Ala Asp Pro Ala Thr Phe  
 225 230 235 240  
 Met Gly Met Ala Gly Leu Gly Asp Leu Val Leu Thr Cys Thr Asp Asn  
 245 250 255  
 Gln Ser Arg Asn Arg Arg Phe Gly Met Met Leu Gly Gln Gly Met Asp  
 260 265 270  
 Val Gln Ser Ala Gln Glu Lys Ile Gly Gln Val Val Glu Gly Tyr Arg  
 275 280 285  
 Asn Thr Lys Glu Val Arg Glu Leu Ala His Arg Phe Gly Val Glu Met  
 290 295 300  
 Pro Ile Thr Glu Glu Ile Tyr Gln Val Leu Tyr Cys Gly Lys Asn Ala  
 305 310 315 320  
 Arg Glu Ala Ala Leu Thr Leu Leu Gly Arg Ala Arg Lys Asp Glu Arg  
 325 330 335  
 Ser Ser His

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 501 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met Glu Thr Lys Asp Leu Ile Val Ile Gly Gly Gly Ile Asn Gly Ala  
 1 5 10 15  
 Gly Ile Ala Ala Asp Ala Ala Gly Arg Gly Leu Ser Val Leu Met Leu  
 20 25 30  
 Glu Ala Gln Asp Leu Ala Cys Ala Thr Ser Ser Ala Ser Ser Lys Leu  
 35 40 45  
 Ile His Gly Gly Leu Arg Tyr Leu Glu His Tyr Glu Phe Arg Leu Val  
 50 55 60

Ser Glu Ala Leu Ala Glu Arg Glu Val Leu Leu Lys Met Ala Pro His  
 65 70 75 80  
 Ile Ala Phe Pro Met Arg Phe Arg Leu Pro His Arg Pro His Leu Arg  
 85 90 95  
 Pro Ala Trp Met Ile Arg Ile Gly Leu Phe Met Tyr Asp His Leu Gly  
 100 105 110  
 Lys Arg Thr Ser Leu Pro Gly Ser Thr Gly Leu Arg Phe Gly Ala Asn  
 115 120 125  
 Ser Val Leu Lys Pro Glu Ile Lys Arg Gly Phe Glu Tyr Ser Asp Cys  
 130 135 140  
 Trp Val Asp Asp Ala Arg Leu Val Leu Ala Asn Ala Gln Met Val Val  
 145 150 155 160  
 Arg Lys Gly Gly Glu Val Leu Thr Arg Thr Arg Ala Thr Ser Ala Arg  
 165 170 175  
 Arg Glu Asn Gly Leu Trp Ile Val Glu Ala Glu Asp Ile Asp Thr Gly  
 180 185 190  
 Lys Lys Tyr Ser Trp Gln Ala Arg Gly Leu Val Asn Ala Thr Gly Pro  
 195 200 205  
 Trp Val Lys Gln Phe Phe Asp Asp Gly Met His Leu Pro Ser Pro Tyr  
 210 215 220  
 Gly Ile Arg Leu Ile Lys Gly Ser His Ile Val Val Pro Arg Val His  
 225 230 235 240  
 Thr Gln Lys Gln Ala Tyr Ile Leu Gln Asn Glu Asp Lys Arg Ile Val  
 245 250 255  
 Phe Val Ile Pro Trp Met Asp Glu Phe Ser Ile Ile Gly Thr Thr Asp  
 260 265 270  
 Val Glu Tyr Lys Gly Asp Pro Lys Ala Val Lys Ile Glu Glu Ser Glu  
 275 280 285  
 Ile Asn Tyr Leu Leu Asn Val Tyr Asn Thr His Phe Lys Lys Gln Leu  
 290 295 300  
 Ser Arg Asp Asp Ile Val Trp Thr Tyr Ser Gly Val Arg Pro Leu Cys  
 305 310 315 320  
 Asp Asp Glu Ser Asp Ser Pro Gln Ala Ile Thr Arg Asp Tyr Thr Leu  
 325 330 335  
 Asp Ile His Asp Glu Asn Gly Lys Ala Pro Leu Leu Ser Val Phe Gly  
 340 345 350  
 Gly Lys Leu Thr Thr Tyr Arg Lys Leu Ala Glu His Ala Leu Glu Lys  
 355 360 365

Leu Thr Pro Tyr Tyr Gln Gly Ile Gly Pro Ala Trp Thr Lys Glu Ser  
 370 375 380  
 Val Leu Pro Gly Gly Ala Ile Glu Gly Asp Arg Asp Asp Tyr Ala Ala  
 385 390 395 400  
 Arg Leu Arg Arg Arg Tyr Pro Phe Leu Thr Glu Ser Leu Ala Arg His  
 405 410 415  
 Tyr Ala Arg Thr Tyr Gly Ser Asn Ser Glu Leu Leu Leu Gly Asn Ala  
 420 425 430  
 Gly Thr Val Ser Asp Leu Gly Glu Asp Phe Gly His Glu Phe Tyr Glu  
 435 440 445  
 Ala Glu Leu Lys Tyr Leu Val Asp His Glu Trp Val Arg Arg Ala Asp  
 450 455 460  
 Asp Ala Leu Trp Arg Arg Thr Lys Gln Gly Met Trp Leu Asn Ala Asp  
 465 470 475 480  
 Gln Gln Ser Arg Val Ser Gln Trp Leu Val Glu Tyr Thr Gln Gln Arg  
 485 490 495  
 Leu Ser Leu Ala Ser  
 500

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 542 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: unknown
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Lys Thr Arg Asp Ser Gln Ser Ser Asp Val Ile Ile Ile Gly Gly  
 1 5 10 15  
 Gly Ala Thr Gly Ala Gly Ile Ala Arg Asp Cys Ala Leu Arg Gly Leu  
 20 25 30  
 Arg Val Ile Leu Val Glu Arg His Asp Ile Ala Thr Gly Ala Thr Gly  
 35 40 45  
 Arg Asn His Gly Leu Leu His Ser Gly Ala Arg Tyr Ala Val Thr Asp  
 50 55 60  
 Ala Glu Ser Ala Arg Glu Cys Ile Ser Glu Asn Gln Ile Leu Lys Arg  
 65 70 75 80  
 Ile Ala Arg His Cys Val Glu Pro Thr Asn Gly Leu Phe Ile Thr Leu  
 85 90 95

Pro Glu Asp Asp Leu Ser Phe Gln Ala Thr Phe Ile Arg Ala Cys Glu  
 100 105 110  
 Glu Ala Gly Ile Ser Ala Glu Ala Ile Asp Pro Gln Gln Ala Arg Ile  
 115 120 125  
 Ile Glu Pro Ala Val Asn Pro Ala Leu Ile Gly Ala Val Lys Val Pro  
 130 135 140  
 Asp Gly Thr Val Asp Pro Phe Arg Leu Thr Ala Ala Asn Met Leu Asp  
 145 150 155 160  
 Ala Lys Glu His Gly Ala Val Ile Leu Thr Ala His Glu Val Thr Gly  
 165 170 175  
 Leu Ile Arg Glu Gly Ala Thr Val Cys Gly Val Arg Val Arg Asn His  
 180 185 190  
 Leu Thr Gly Glu Thr Gln Ala Leu His Ala Pro Val Val Val Asn Ala  
 195 200 205  
 Ala Gly Ile Trp Gly Gln His Ile Ala Glu Tyr Ala Asp Leu Arg Ile  
 210 215 220  
 Arg Met Phe Pro Ala Lys Gly Ser Leu Leu Ile Met Asp His Arg Ile  
 225 230 235 240  
 Asn Gln His Val Ile Asn Arg Cys Arg Lys Pro Ser Asp Ala Asp Ile  
 245 250 255  
 Leu Val Pro Gly Asp Thr Ile Ser Leu Ile Gly Thr Thr Ser Leu Arg  
 260 265 270  
 Ile Asp Tyr Asn Glu Ile Asp Asp Asn Arg Val Thr Ala Glu Glu Val  
 275 280 285  
 Asp Ile Leu Leu Arg Glu Gly Glu Lys Leu Ala Pro Val Met Ala Lys  
 290 295 300  
 Thr Arg Ile Leu Arg Ala Tyr Ser Gly Val Arg Pro Leu Val Ala Ser  
 305 310 315 320  
 Asp Asp Asp Pro Ser Gly Arg Asn Leu Ser Arg Gly Ile Val Leu Leu  
 325 330 335  
 Asp His Ala Glu Arg Asp Gly Leu Asp Gly Phe Ile Thr Ile Thr Gly  
 340 345 350  
 Gly Lys Leu Met Thr Tyr Arg Leu Met Ala Glu Trp Ala Thr Asp Ala  
 355 360 365  
 Val Cys Arg Lys Leu Gly Asn Thr Arg Pro Cys Thr Thr Ala Asp Leu  
 370 375 380  
 Ala Leu Pro Gly Ser Gln Glu Pro Ala Glu Val Thr Leu Arg Lys Val  
 385 390 395 400

Ile Ser Leu Pro Ala Pro Leu Arg Gly Ser Ala Val Tyr Arg His Gly  
 405 410 415  
 Asp Arg Thr Pro Ala Trp Leu Ser Glu Gly Arg Leu His Arg Ser Leu  
 420 425 430  
 Val Cys Glu Cys Glu Ala Val Thr Ala Gly Glu Val Gln Tyr Ala Val  
 435 440 445  
 Glu Asn Leu Asn Val Asn Ser Leu Leu Asp Leu Arg Arg Arg Thr Arg  
 450 455 460  
 Val Gly Met Gly Thr Cys Gln Gly Glu Leu Cys Ala Cys Arg Ala Ala  
 465 470 475 480  
 Gly Leu Leu Gln Arg Phe Asn Val Thr Thr Ser Ala Gln Ser Ile Glu  
 485 490 495  
 Gln Leu Ser Thr Phe Leu Asn Glu Arg Trp Lys Gly Val Gln Pro Ile  
 500 505 510  
 Ala Trp Gly Asp Ala Leu Arg Glu Ser Glu Phe Thr Arg Trp Val Tyr  
 515 520 525  
 Gln Gly Leu Cys Gly Leu Glu Lys Glu Gln Lys Asp Ala Leu  
 530 535 540

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met Gly Leu Thr Thr Lys Pro Leu Ser Leu Lys Val Asn Ala Ala Leu  
 1 5 10 15  
 Phe Asp Val Asp Gly Thr Ile Ile Ile Ser Gln Pro Ala Ile Ala Ala  
 20 25 30  
 Phe Trp Arg Asp Phe Gly Lys Asp Lys Pro Tyr Phe Asp Ala Glu His  
 35 40 45  
 Val Ile Gln Val Ser His Gly Trp Arg Thr Phe Asp Ala Ile Ala Lys  
 50 55 60  
 Phe Ala Pro Asp Phe Ala Asn Glu Glu Tyr Val Asn Lys Leu Glu Ala  
 65 70 75 80  
 Glu Ile Pro Val Lys Tyr Gly Glu Lys Ser Ile Glu Val Pro Gly Ala  
 85 90 95



Val Lys Leu Cys Asn Ala Leu Asn Ala Leu Pro Lys Glu Lys Trp Ala  
 100 105 110  
 Val Ala Thr Ser Gly Thr Arg Asp Met Ala Gln Lys Trp Phe Glu His  
 115 120 125  
 Leu Gly Ile Arg Arg Pro Lys Tyr Phe Ile Thr Ala Asn Asp Val Lys  
 130 135 140  
 Gln Gly Lys Pro His Pro Glu Pro Tyr Leu Lys Gly Arg Asn Gly Leu  
 145 150 155 160  
 Gly Tyr Pro Ile Asn Glu Gln Asp Pro Ser Lys Ser Lys Val Val Val  
 165 170 175  
 Phe Glu Asp Ala Pro Ala Gly Ile Ala Ala Gly Lys Ala Ala Gly Cys  
 180 185 190  
 Lys Ile Ile Gly Ile Ala Thr Thr Phe Asp Leu Asp Phe Leu Lys Glu  
 195 200 205  
 Lys Gly Cys Asp Ile Ile Val Lys Asn His Glu Ser Ile Arg Val Gly  
 210 215 220  
 Gly Tyr Asn Ala Glu Thr Asp Glu Val Glu Phe Ile Phe Asp Asp Tyr  
 225 230 235 240  
 Leu Tyr Ala Lys Asp Asp Leu Leu Lys Trp  
 245 250

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 271 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Lys Arg Phe Asn Val Leu Lys Tyr Ile Arg Thr Thr Lys Ala Asn  
 1 5 10 15  
 Ile Gln Thr Ile Ala Met Pro Leu Thr Thr Lys Pro Leu Ser Leu Lys  
 20 25 30  
 Ile Asn Ala Ala Leu Phe Asp Val Asp Gly Thr Ile Ile Ile Ser Gln  
 35 40 45  
 Pro Ala Ile Ala Ala Phe Trp Arg Asp Phe Gly Lys Asp Lys Pro Tyr  
 50 55 60  
 Phe Asp Ala Glu His Val Ile His Ile Ser His Gly Trp Arg Thr Tyr  
 65 70 75 80

Asp Ala Ile Ala Lys Phe Ala Pro Asp Phe Ala Asp Glu Glu Tyr Val  
 85 90 95  
 Asn Lys Leu Glu Gly Glu Ile Pro Glu Lys Tyr Gly Glu His Ser Ile  
 100 105 110  
 Glu Val Pro Gly Ala Val Lys Leu Cys Asn Ala Leu Asn Ala Leu Pro  
 115 120 125  
 Lys Glu Lys Trp Ala Val Ala Thr Ser Gly Thr Arg Asp Met Ala Lys  
 130 135 140  
 Lys Trp Phe Asp Ile Leu Lys Ile Lys Arg Pro Glu Tyr Phe Ile Thr  
 145 150 155 160  
 Ala Asn Asp Val Lys Gln Gly Lys Pro His Pro Glu Pro Tyr Leu Lys  
 165 170 175  
 Gly Arg Asn Gly Leu Gly Phe Pro Ile Asn Glu Gln Asp Pro Ser Lys  
 180 185 190  
 Ser Lys Val Val Val Phe Glu Asp Ala Pro Ala Gly Ile Ala Ala Gly  
 195 200 205  
 Lys Ala Ala Gly Cys Lys Ile Val Gly Ile Ala Thr Thr Phe Asp Leu  
 210 215 220  
 Asp Phe Leu Lys Glu Lys Gly Cys Asp Ile Ile Val Lys Asn His Glu  
 225 230 235 240  
 Ser Ile Arg Val Gly Glu Tyr Asn Ala Glu Thr Asp Glu Val Glu Leu  
 245 250 255  
 Ile Phe Asp Asp Tyr Leu Tyr Ala Lys Asp Asp Leu Leu Lys Trp  
 260 265 270

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 709 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: unknown
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Met Phe Pro Ser Leu Phe Arg Leu Val Val Phe Ser Lys Arg Tyr Ile  
 1 5 10 15  
 Phe Arg Ser Ser Gln Arg Leu Tyr Thr Ser Leu Lys Gln Glu Gln Ser  
 20 25 30  
 Arg Met Ser Lys Ile Met Glu Asp Leu Arg Ser Asp Tyr Val Pro Leu  
 35 40 45

Ile	Ala	Ser	Ile	Asp	Val	Gly	Thr	Thr	Ser	Ser	Arg	Cys	Ile	Leu	Phe	50	55	60
Asn	Arg	Trp	Gly	Gln	Asp	Val	Ser	Lys	His	Gln	Ile	Glu	Tyr	Ser	Thr	65	70	75
Ser	Ala	Ser	Lys	Gly	Lys	Ile	Gly	Val	Ser	Gly	Leu	Arg	Arg	Pro	Ser	85	90	95
Thr	Ala	Pro	Ala	Arg	Glu	Thr	Pro	Asn	Ala	Gly	Asp	Ile	Lys	Thr	Ser	100	105	110
Gly	Lys	Pro	Ile	Phe	Ser	Ala	Glu	Gly	Tyr	Ala	Ile	Gln	Glu	Thr	Lys	115	120	125
Phe	Leu	Lys	Ile	Glu	Glu	Leu	Asp	Leu	Asp	Phe	His	Asn	Glu	Pro	Thr	130	135	140
Leu	Lys	Phe	Pro	Lys	Pro	Gly	Trp	Val	Glu	Cys	His	Pro	Gln	Lys	Leu	145	150	155
Leu	Val	Asn	Val	Val	Gln	Cys	Leu	Ala	Ser	Ser	Leu	Leu	Ser	Leu	Gln	165	170	175
Thr	Ile	Asn	Ser	Glu	Arg	Val	Ala	Asn	Gly	Leu	Pro	Pro	Tyr	Lys	Val	180	185	190
Ile	Cys	Met	Gly	Ile	Ala	Asn	Met	Arg	Glu	Thr	Thr	Ile	Leu	Trp	Ser	195	200	205
Arg	Arg	Thr	Gly	Lys	Pro	Ile	Val	Asn	Tyr	Gly	Ile	Val	Trp	Asn	Asp	210	215	220
Thr	Arg	Thr	Ile	Lys	Ile	Val	Arg	Asp	Lys	Trp	Gln	Asn	Thr	Ser	Val	225	230	235
Asp	Arg	Gln	Leu	Gln	Leu	Arg	Gln	Lys	Thr	Gly	Leu	Pro	Leu	Leu	Ser	245	250	255
Thr	Tyr	Phe	Ser	Cys	Ser	Lys	Leu	Arg	Trp	Phe	Leu	Asp	Asn	Glu	Pro	260	265	270
Leu	Cys	Thr	Lys	Ala	Tyr	Glu	Glu	Asn	Asp	Leu	Met	Phe	Gly	Thr	Val	275	280	285
Asp	Thr	Trp	Leu	Ile	Tyr	Gln	Leu	Thr	Lys	Gln	Lys	Ala	Phe	Val	Ser	290	295	300
Asp	Val	Thr	Asn	Ala	Ser	Arg	Thr	Gly	Phe	Met	Asn	Leu	Ser	Thr	Leu	305	310	315
Lys	Tyr	Asp	Asn	Glu	Leu	Leu	Glu	Phe	Trp	Gly	Ile	Asp	Lys	Asn	Leu	325	330	335
Ile	His	Met	Pro	Glu	Ile	Val	Ser	Ser	Ser	Gln	Tyr	Tyr	Gly	Asp	Phe	340	345	350

Gly Ile Pro Asp Trp Ile Met Glu Lys Leu His Asp Ser Pro Lys Thr  
 355 360 365  
 Val Leu Arg Asp Leu Val Lys Arg Asn Leu Pro Ile Gln Gly Cys Leu  
 370 375 380  
 Gly Asp Gln Ser Ala Ser Met Val Gly Gln Leu Ala Tyr Lys Pro Gly  
 385 390 395 400  
 Ala Ala Lys Cys Thr Tyr Gly Thr Gly Cys Phe Leu Leu Tyr Asn Thr  
 405 410 415  
 Gly Thr Lys Lys Leu Ile Ser Gln His Gly Ala Leu Thr Thr Leu Ala  
 420 425 430  
 Phe Trp Phe Pro His Leu Gln Glu Tyr Gly Gly Gln Lys Pro Glu Leu  
 435 440 445  
 Ser Lys Pro His Phe Ala Leu Glu Gly Ser Val Ala Val Ala Gly Ala  
 450 455 460  
 Val Val Gln Trp Leu Arg Asp Asn Leu Arg Leu Ile Asp Lys Ser Glu  
 465 470 475 480  
 Asp Val Gly Pro Ile Ala Ser Thr Val Pro Asp Ser Gly Gly Val Val  
 485 490 495  
 Phe Val Pro Ala Phe Ser Gly Leu Phe Ala Pro Tyr Trp Asp Pro Asp  
 500 505 510  
 Ala Arg Ala Thr Ile Met Gly Met Ser Gln Phe Thr Thr Ala Ser His  
 515 520 525  
 Ile Ala Arg Ala Ala Val Glu Gly Val Cys Phe Gln Ala Arg Ala Ile  
 530 535 540  
 Leu Lys Ala Met Ser Ser Asp Ala Phe Gly Glu Gly Ser Lys Asp Arg  
 545 550 555 560  
 Asp Phe Leu Glu Glu Ile Ser Asp Val Thr Tyr Glu Lys Ser Pro Leu  
 565 570 575  
 Ser Val Leu Ala Val Asp Gly Gly Met Ser Arg Ser Asn Glu Val Met  
 580 585 590  
 Gln Ile Gln Ala Asp Ile Leu Gly Pro Cys Val Lys Val Arg Arg Ser  
 595 600 605  
 Pro Thr Ala Glu Cys Thr Ala Leu Gly Ala Ala Ile Ala Ala Asn Met  
 610 615 620  
 Ala Phe Lys Asp Val Asn Glu Arg Pro Leu Trp Lys Asp Leu His Asp  
 625 630 635 640  
 Val Lys Lys Trp Val Phe Tyr Asn Gly Met Glu Lys Asn Glu Gln Ile  
 645 650 655

Ser Pro Glu Ala His Pro Asn Leu Lys Ile Phe Arg Ser Glu Ser Asp  
                   660                                  665                                  670

Asp Ala Glu Arg Arg Lys His Trp Lys Tyr Trp Glu Val Ala Val Glu  
                   675                                  680                                  685

Arg Ser Lys Gly Trp Leu Lys Asp Ile Glu Gly Glu His Glu Gln Val  
                   690                                  695                                  700

Leu Glu Asn Phe Gln  
                   705

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 51 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GCGCGGATCC AGGAGTCTAG AATTATGGGA TTGACTACTA AACCTCTATC T 51

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GATACGCCCCG GGTTACCATT TCAACAGATC GTCCTT 36

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 34 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TTGATAATAT AACCATGGCT GCTGCTGCTG ATAG 34

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 39 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GTATGATATG TTATCTTGGA TCCAATAAAT CTAATCTTC

39

## (2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CATGACTAGT AAGGAGGACA ATTC

24

## (2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CATGGAATTG TCCTCCTTAC TAGT

24

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CTAGTAAGGA GGACAATTC

19



## (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CATGGAATTG TCCTCCTTA

19

## (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 15 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (iii) HYPOTHETICAL: NO

- (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GATCCAGGAA ACAGA

15

## (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 15 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (iii) HYPOTHETICAL: NO

- (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CTAGTCTGTT TCCTG

15

## (2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GCTTTCTGTG CTGCGGCTTT AG

22

- (2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TGGTCGAGGA TCCACTTCAC TTT

23

- (2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 51 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

AAAGTGAAGT GGATCCTCGA CCAATTGGAT GGTGGCGCAG TAGCAAACAA T

51

- (2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GGATCACCGC CGCAGAACT ACG

23

- (2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
  - (A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:  
CTGTCAGCCG TTAAGTGTTT CTGTG 25
- (2) INFORMATION FOR SEQ ID NO:31:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:  
CAGTTCAACC TGTGATAGT ACG 23
- (2) INFORMATION FOR SEQ ID NO:32:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:  
ATGAGTCAAA CATCAACCTT 20
- (2) INFORMATION FOR SEQ ID NO:33:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:  
ATGGAGAAAA AAATCACTGG 20
- (2) INFORMATION FOR SEQ ID NO:34:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:  
TTACGCCCCG CCCTGCCACT 20

## (2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TCAGAGGATG TGCACCTGCA

20

## (2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

CGAGCATGCC GCATTTGGCA CTACTC

26

## (2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GCGTCTAGAG TAGGTTATTC CCACTCTTG

29

## (2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GAAGTCGACC GCTGCGCCTT ATCCGG

26

## (2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

CGCGTCGACG TTTACAATTT CAGGTGGC

28

## (2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GCAGCATGCT GGACTGGTAG TAG

23

## (2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

CAGTCTAGAG TTATTGGCAA ACCTACC

27

## (2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GATGCATGCC CAGGGCGGAG ACGGC

25

## (2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PRIMER"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CTAACGATTG TTCTCTAGAG AAAATGTCC

29

## (2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "primer"

- (iii) HYPOTHETICAL: NO

- (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

CACGCATGCA GTTCAACCTG TTGATAGTAC

30

## (2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "primer"

- (iii) HYPOTHETICAL: NO

- (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GCGTCTAGAT CCTTTTAAAT TAAAAATG

28



# INTERNATIONAL SEARCH REPORT

Inte. onal Application No  
PCT/US 98/25551

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6    C12N15/53    C12N15/55    C12N15/60    C12P7/20    C12P7/18 C12N1/15    C12N1/19    C12N1/21    C12N9/04    C12N9/16 C12N9/88		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6    C12N    C12P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WANG H.-T. ET AL.: "Cloning, sequence, and disruption of Saccharomyces diastaticus DAR1 gene encoding a glycerol-3-phosphate dehydrogenase." JOURNAL OF BACTERIOLOGY, vol. 176, no. 22, November 1994, pages 7091-7095, XP000563880 cited in the application see abstract see page 7091, column 2, paragraph 2 <div style="text-align: center;">--- -/--</div>	1-16
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
° Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  <div style="text-align: center; font-weight: bold;">5 March 1999</div>		Date of mailing of the international search report  <div style="text-align: center; font-weight: bold;">12/03/1999</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  <div style="text-align: center; font-weight: bold;">Hornig, H</div>

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/25551

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	NORBECK J. ET AL.: "Purification and characterization of two isoenzymes of DL-glycerol-3-phosphatase from <i>Saccharomyces cerevisiae</i> ." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 23, 7 June 1996, pages 13875-13881, XP002058248 cited in the application see abstract see page 13875, column 2, paragraph 2 see page 13881, column 1, line 8 - line 21 ---	1-16
A	HIRAYAMA T ET AL: "CLONING AND CHARACTERIZATION OF SEVEN CDNAS FOR HYPEROSMOLARITY- RESPONSIVE (HOR) GENES OF <i>SACCHAROMYCES CEREVISIAE</i> " MOLECULAR AND GENERAL GENETICS, vol. 249, 1995, pages 127-138, XP002058249 cited in the application see abstract see page 129, right-hand column, line 2; figure 1A ---	1-16
A	LARSSON K. ET AL.: "A gene encoding sn-glycerol 3-phosphate dehydrogenase (NAD+) complements an osmosensitive mutant of <i>Saccharomyces cerevisiae</i> ." MOLECULAR MICROBIOLOGY, vol. 10, no. 5, 1993, pages 1101-1111, XP000562759 cited in the application see abstract ---	1-16
A	FR 2 735 145 A (AGRONOMIQUE INST NAT RECH) 13 December 1996 see claims 1-5 ---	1-16
A	WO 97 07199 A (WISCONSIN ALUMNI RES FOUND) 27 February 1997 see claims 9-12 ---	1-16
A	OMORI T. ET AL.: "Breeding of high glycerol-producing shochu yeast ( <i>Saccharomyces cerevisiae</i> ) with acquired salt tolerance." JOURNAL OF FERMENTATION AND BIOENGINEERING, vol. 79, no. 6, 1995, pages 560-565, XP002058250 see abstract ---	1-16
P,A	WO 98 21340 A (BULTHUIS BEN A ;HAYNIE SHARON LORETTA (US); HSU AMY KUANG HUA (US)) 22 May 1998 see the whole document ---	1-16
	-/--	

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/25551

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 35795 A (DU PONT ;NAGARAJAN VASANTHA (US); NAKAMURA CHARLES EDWIN (US)) 14 November 1996 see the whole document ---	17-20
A	WO 96 35796 A (DU PONT ;GENENCOR INT (US); LAFFEND LISA ANNE (US); NAGARAJAN VASA) 14 November 1996 see the whole document ---	17-20
A	WO 96 35799 A (DU PONT ;HAYNIE SHARON LORETTA (US); WAGNER LORRAINE WINONA (US)) 14 November 1996 see the whole document ---	17-20
P,A	WO 98 21339 A (DIAS TORRES MARIA ;HAYNIE SHARON LORETTA (US); HSU AMY KUANG HUA ( ) 22 May 1998 see the whole document ---	17-20
P,A	WO 98 21341 A (DIAZ TORRES MARIA ;CHASE MATTHEW W (US); GENENCOR INT (US); TRIMBU) 22 May 1998 see the whole document -----	17-20

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/US 98/25551

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2735145 A	13-12-1996	AU 6229796 A WO 9641888 A	09-01-1997 27-12-1996
WO 9707199 A	27-02-1997	US 5733749 A AU 6715196 A CA 2229488 A EP 0845030 A	31-03-1998 12-03-1997 27-02-1997 03-06-1998
WO 9821340 A	22-05-1998	AU 5430798 A	03-06-1998
WO 9635795 A	14-11-1996	US 5633362 A AU 5722996 A CA 2220441 A EP 0827543 A US 5821092 A	27-05-1997 29-11-1996 14-11-1996 11-03-1998 13-10-1998
WO 9635796 A	14-11-1996	US 5686276 A AU 5678996 A CA 2220880 A EP 0826057 A	11-11-1997 29-11-1996 14-11-1996 04-03-1998
WO 9635799 A	14-11-1996	US 5599689 A AU 5722896 A CA 2220396 A EP 0824595 A JP 10507082 T	04-02-1997 29-11-1996 14-11-1996 25-02-1998 14-07-1998
WO 9821339 A	22-05-1998	AU 5248498 A AU 5507698 A WO 9821341 A	03-06-1998 03-06-1998 22-05-1998
WO 9821341 A	22-05-1998	AU 5248498 A AU 5507698 A WO 9821339 A	03-06-1998 03-06-1998 22-05-1998